

Consistency of Heckman-type two-step Estimators for the Multivariate Sample-Selection Model

Tauchmann, Harald

Postprint / Postprint

Zeitschriftenartikel / journal article

Zur Verfügung gestellt in Kooperation mit / provided in cooperation with:

www.peerproject.eu

Empfohlene Zitierung / Suggested Citation:

Tauchmann, H. (2009). Consistency of Heckman-type two-step Estimators for the Multivariate Sample-Selection Model. *Applied Economics*, 42(30), 3895-. <https://doi.org/10.1080/00036840802360179>

Nutzungsbedingungen:

Dieser Text wird unter dem "PEER Licence Agreement zur Verfügung" gestellt. Nähere Auskünfte zum PEER-Projekt finden Sie hier: <http://www.peerproject.eu> Gewährt wird ein nicht exklusives, nicht übertragbares, persönliches und beschränktes Recht auf Nutzung dieses Dokuments. Dieses Dokument ist ausschließlich für den persönlichen, nicht-kommerziellen Gebrauch bestimmt. Auf sämtlichen Kopien dieses Dokuments müssen alle Urheberrechtshinweise und sonstigen Hinweise auf gesetzlichen Schutz beibehalten werden. Sie dürfen dieses Dokument nicht in irgendeiner Weise abändern, noch dürfen Sie dieses Dokument für öffentliche oder kommerzielle Zwecke vervielfältigen, öffentlich ausstellen, aufführen, vertreiben oder anderweitig nutzen.

Mit der Verwendung dieses Dokuments erkennen Sie die Nutzungsbedingungen an.

Terms of use:

This document is made available under the "PEER Licence Agreement". For more information regarding the PEER-project see: <http://www.peerproject.eu> This document is solely intended for your personal, non-commercial use. All of the copies of this documents must retain all copyright information and other information regarding legal protection. You are not allowed to alter this document in any way, to copy it for public or commercial purposes, to exhibit the document in public, to perform, distribute or otherwise use the document in public.

By using this particular document, you accept the above-stated conditions of use.



Consistency of Heckman-type two-step Estimators for the Multivariate Sample-Selection Model

Journal:	<i>Applied Economics</i>
Manuscript ID:	APE-06-0698
Journal Selection:	Applied Economics
JEL Code:	C15 - Statistical Simulation Methods Monte Carlo Methods < C1 - Econometric and Statistical Methods: General < C - Mathematical and Quantitative Methods, C34 - Truncated and Censored Models < C3 - Econometric Methods: Multiple/Simultaneous Equation Models < C - Mathematical and Quantitative Methods, C51 - Model Construction and Estimation < C5 - Econometric Modeling < C - Mathematical and Quantitative Methods
Keywords:	multivariate sample-selection model, censored system of equations, Heckman-correction



Consistency of Heckman-type two-step Estimators for the Multivariate Sample-Selection Model

November 2006

Abstract: This analysis shows that multivariate generalizations to the classical Heckman (1976 and 1979) two-step estimator that account for cross-equation correlation and use the inverse Mills ratio as correction-term are consistent only if certain restrictions apply to the true error-covariance structure. An alternative class of generalizations to the classical Heckman two-step approach is derived that condition on the entire selection pattern rather than selection in particular equations and, therefore, use modified correction-terms. It is shown that this class of estimators is consistent. In addition, Monte-Carlo results illustrate that these estimators display a smaller mean square prediction error.

JEL Classification Number: C15, C34, C51.

Key words: multivariate sample-selection model, censored system of equations, Heckman-correction.

1 Introduction

Using non-aggregated micro-data for estimating systems of seemingly unrelated equations – the most prominent among them being demand systems – often encounters the problem of numerous zero-observations in the dependent variables. These cannot be appropriately explained by conventional continuous *SUR*¹ models. Instead, zero-observations may be modelled as determined by an upstream multivariate binary choice problem. Under the assumption of normally distributed errors, the resulting joint model represents a multivariate generalization to the classical univariate sample-selection model, cf. Heckman (1976 and 1979). In the literature, this model is often referred to as a “censored system of equations”, yet censoring in the narrow sense just represents a special case of the general model.²

The question of how to estimate the parameters of this model is subject to an ongoing debate. Clearly, under parametric distributional assumptions full information maximum likelihood (*FIML*) is the efficient estimation technique. In fact, *FIML* has recently been applied to this problem by Yen (2005). However, the *FIML* estimator is computationally extremely demanding, rendering much simpler two-step approaches worth considering for many applications.

Among two-step estimators the one proposed by Heien & Wessels (1990) has been particularly popular. Besides numerous other authors, it has been applied by Heien

¹See Zellner (1963) for the seemingly unrelated regression equations (*SUR*) model.

²We stick to the relevant literature and use the term “censored” as a synonym for “not selected”.

1
2
3
4 & Durham (1991), Gao et al. (1995), and Nayga et al. (1999). However, Shonkwiler
5
6 & Yen (1999) as well as Vermeulen (2001) show that this estimator lacks a decent
7
8 basis in statistical theory and cannot be interpreted in terms of conditional means.
9
10 The Heien & Wessels (1990) estimator, therefore, is inconsistent despite its popularity.
11
12 Chen & Yen (2005) further investigate the nature of its inconsistency and show that
13
14 even a modified variant of this estimator fails to correct properly for sample-selection
15
16 bias. Shonkwiler & Yen (1999) propose an alternative simple two-step estimator that
17
18 – in contrast to Heien & Wessels (1990) – is theoretically well founded. This estimator
19
20 is based on the mean of dependent variables that is unconditional on the outcome of
21
22 the upstream discrete choice model. Su & Yen (2000), Yen et al. (2002) and Goodwin
23
24 et al. (2004) may serve as examples for applications of this procedure.
25
26
27
28
29
30

31
32 Tauchmann (2005) compares the performance of the Shonkwiler & Yen (1999) esti-
33
34 mator and two-step estimators that – analogously to the classical Heckman (1976 and
35
36 1979) two-step approach, yet in contrast to Shonkwiler & Yen (1999) – condition on
37
38 the outcome the upstream discrete choice model. In terms of the mean square predic-
39
40 tion error, the unconditional Shonkwiler & Yen (1999) estimator is shown to perform
41
42 poorly if the conditional mean of the dependent variables is large compared to its
43
44 conditional variance. Tauchmann (2005), however, exclusively focuses on the mean
45
46 square error yet does not check for unbiasedness and consistency of the conditional
47
48 estimators. Though one may argue that it is of no relevance in applied work whether
49
50 an error originates from an estimator's bias or from its variance, many researches do
51
52
53
54
55
56
57
58
59
60

1
2
3
4 avoid inconsistent estimators, even if their mean square error is small. For this rea-
5
6
7 son, addressing unbiasedness and consistency of Heckman-type two-step estimators for
8
9 censored systems of equations is a relevant task.
10

11
12 The analysis presented in this article shows that some of the estimators proposed
13
14 by Tauchmann (2005) are consistent only for restrictive error-covariance structures. It
15
16 also shows that a modified two-step Heckman-type estimator is generally consistent
17
18 and performs well in terms of the mean square prediction error. In order to yield
19
20 these results, the remainder of this paper is organized as follows: Section 2 introduces
21
22 the model to be analyzed in more detail and analyzes the properties of straightfor-
23
24 ward multivariate generalizations to the Heckman (1976 and 1979) two-step estimator.
25
26 In Section 3 an alternative class of generalized two-step Heckman-type estimators is
27
28 derived. Section 4 presents results from Monte-Carlo simulations that illustrate the
29
30 theoretical results and extends the analysis to the estimators' mean square error. Sec-
31
32 tion 5 concludes.
33
34
35
36
37
38
39
40
41

42 **2 An analysis of sample-selection models**

43 44 45 **2.1 A multivariate sample-selection model**

46
47
48 Recall the m -variate sample-selection model, which is analyzed by Heinen & Wessels
49
50 (1990), Shonkwiler & Yen (1999), Tauchmann (2005), Yen (2005), and Chen & Yen
51
52
53
54
55
56
57
58
59
60

(2005). The equations

$$y_{it}^* = x_{it}'\beta_i + \varepsilon_{it} \quad (1)$$

$$d_{it}^* = z_{it}'\alpha_i + v_{it}, \quad (2)$$

characterize the latent model, that is y_{it}^* and d_{it}^* are unobserved. Their observed counterparts y_{it} and d_{it} are determined by

$$d_{it} = \begin{cases} 1 & \text{if } d_{it}^* > 0 \\ 0 & \text{if } d_{it}^* \leq 0 \end{cases} \quad (3)$$

$$y_{it} = d_{it}y_{it}^*. \quad (4)$$

Here, $i = 1, \dots, m$ indexes the m equations of the system, and $t = 1, \dots, T$ indexes the individuals. x_{it} and z_{it} are vectors of observed exogenous variables. The vector $d_t = [d_{1t} \dots d_{mt}]'$ describes the entire individual selection pattern. Finally, $\varepsilon_t = [\varepsilon_{1t} \dots \varepsilon_{mt}]'$ and $v_t = [v_{1t} \dots v_{mt}]'$ are normally distributed, zero-mean error vectors with the covariance matrix

$$\text{Var}(\varepsilon_t, v_t) = \begin{bmatrix} \Sigma_{\varepsilon\varepsilon} & \Sigma'_{\varepsilon v} \\ \Sigma_{\varepsilon v} & \Sigma_{vv} \end{bmatrix}. \quad (5)$$

The diagonal-elements of Σ_{vv} are subject to the normalization $\sigma_{ii}^{vv} = 1$, $i = 1 \dots m$.

2.2 Inconsistency of Heckman-type estimators

For the model (1) through (4) Tauchmann (2005) suggests a class of system two-step estimators that – analogously to the original Heckman two-step approach – con-

ditions on d_{it} equation-by-equation. That is, after first-step estimation of the vectors α_i by univariate or multivariate probit, the second-step regressions yielding estimates for the vectors β_i are based on the conditional expectations³ $E(y_{it}|x_{it}, d_{it}) = d_{it}x'_{it}\beta_i + d_{it}\sigma_{ii}^{\varepsilon v}\lambda(z'_{it}\alpha_i)$. Each regression equation, therefore, includes the inverse Mills ratio $\lambda(z'_{it}\hat{\alpha}_i)$ as an auxiliary regressor and the parameters $\sigma_{ii}^{\varepsilon v}$ are estimated as regression coefficients. Note that d_{it} serves as a weighting variable, i.e. censored observations are weighted by zero and are therefore effectively excluded from the regression.⁴

Tauchmann (2005) distinguishes three variants of this estimator: The first one uses ordinary least squares (*OLS*) and ignores cross-equation correlation of ε_{it} , another variant accounts for it in a simplified *SUR* fashion, and a third accounts for cross-equation correlation and heteroscedasticity using a proper generalized least squares (*GLS*) approach.⁵

In order to analyze these estimators' properties, we consider α as known and focus

³To simplify notation, $E(y_{it}|x_{it}, d_{it} = 1)$ is used as short term for $E(y_{it}|x_{it}, v_{it} > -z'_{it}\alpha_i)$ throughout this paper. Yet, it does *not* denote $E_z[E(y_{it}|x_{it}, v_{it} > -z'_{it}\alpha_i)]$, although z_{it} is not explicitly mentioned in list of the conditioning variables. This analogously applies to any moment that is conditional on either $d_{it} = 1$, $d_{it}d_{jt} = 1$, d_{it} , $d_{it}d_{jt}$, or d_t .

⁴Because of (4), which implies $E(y_{it}|x_{it}, d_{it} = 0) = 0$, the original Heckman (1976 and 1979) estimator can well be interpreted as a procedure that conditions on d_{it} in the full sample and, therefore, uses d_{it} as a weighting variable rather than an estimation procedure that conditions on $d_{it} = 1$ and uses the sub-sample of selected units; see Tauchmann (2005) for details.

⁵Because of $\text{var}(\varepsilon_{it}|d_{it} = 1) = \sigma_{ii}^{\varepsilon\varepsilon}((1 - \sigma_{ii}^{\varepsilon v 2}\sigma_{ii}^{\varepsilon\varepsilon-1}) + \sigma_{ii}^{\varepsilon v 2}\sigma_{ii}^{\varepsilon\varepsilon-1}(1 - z'_{it}\alpha_i\lambda(z'_{it}\alpha_i) - \lambda(z'_{it}\alpha_i)^2))$, cf. Heckman (1976), the errors are heteroscedastic and *SUR* is not a proper *GLS* estimator.

on the second-step regression. Let X denote the stacked, $mT \times mk$ regressor-matrix⁶ consisting of rows $[\mathbf{0}_{1 \times k(i-1)} \ x'_{it} \ \lambda(z'_{it}\alpha_i) \ \mathbf{0}_{1 \times k(m-i)}]$. Note that inverse Mills ratios are included to the list of regressors. Let D denote a $mT \times mT$ matrix with diagonal-elements d_{it} and zero off-diagonal elements. This matrix allocates zero weight to censored units. Ω denotes the $mT \times mT$ block-diagonal weighting-matrix with elements ω_{ijt} . It coincides with the identity-matrix if the model is estimated using the classical Heckman approach equation-by-equation, i.e. *OLS*. In the case of *SUR* estimation, the individual $m \times m$ sub-matrices Ω_t are uniform across all t . In the case of *GLS* estimation, these weighting matrices are individually derived through matrix-inversion from estimates for $\text{var}(\varepsilon_{it}|d_{it} = 1)$ and $\text{cov}(\varepsilon_{it}, \varepsilon_{jt}|d_{it}d_{jt} = 1)$. Finally, let Y denote the stacked $mT \times 1$ vector of dependent variables y_{it} and $\tilde{\varepsilon}$ denote the corresponding $mT \times 1$ error-vector. Because of the inclusion of $\lambda(z'_{it}\alpha_i)$ to the list of regressors and $E(\varepsilon_{it}|d_{it} = 1) = \sigma_{ii}^{\varepsilon} \lambda(z'_{it}\alpha_i)$, the error vector $\tilde{\varepsilon}$ consists of elements $\varepsilon_{it} - E(\varepsilon_{it}|d_{it} = 1)$ rather than ε_{it} . Now the generalized Heckman-estimators for $\hat{\beta}$ proposed by Tauchmann (2005) can be written

$$\hat{\beta} = (X'D\Omega DX)^{-1} X'D\Omega Y. \quad (6)$$

Because of $Y = D(X\beta + \tilde{\varepsilon})$ equation (6) is equivalent to

$$\hat{\beta} = \beta + (X'D\Omega DX)^{-1} X'\Xi, \quad \text{with } \Xi \equiv D\Omega D\tilde{\varepsilon}. \quad (7)$$

⁶ k_i denotes the number of coefficients in the i th equation. In order to simplify notation, yet with no loss of generality, we assume $k_i = k$ for $i = 1, \dots, m$. The matrix X is arranged as such that all m rows belonging to an individual t adjoin each other.

Here, the condition $E(\Xi|X) = \mathbf{0}$ implies $\text{plim } T^{-1}(X'\Xi) = \mathbf{0}$ and, therefore, implies consistency of $\widehat{\beta}$ under standard regularity conditions. To check whether $E(\Xi|X) = \mathbf{0}$ holds, consider an arbitrary element from Ξ :

$$\begin{aligned}\xi_{it} &= \omega_{iit}d_{it}\tilde{\varepsilon}_{it} + \sum_{\substack{j=1 \\ j \neq i}}^m \omega_{ijt}d_{it}d_{jt}\tilde{\varepsilon}_{jt} \\ &= \omega_{iit}d_{it}[\varepsilon_{it} - E(\varepsilon_{it}|d_{it} = 1)] + \sum_{\substack{j=1 \\ j \neq i}}^m \omega_{ijt}d_{it}d_{jt}[\varepsilon_{jt} - E(\varepsilon_{jt}|d_{jt} = 1)].\end{aligned}\quad (8)$$

We apply the law of iterated expectations to (8). First, we take the expectation of ξ_{it} conditional on x_t as well as on the individual selection pattern d_t .

$$E(\xi_{it}|x_t, d_t) = \omega_{iit}d_{it}[E(\varepsilon_{it}|d_t) - E(\varepsilon_{it}|d_{it} = 1)] + \sum_{\substack{j=1 \\ j \neq i}}^m \omega_{ijt}d_{it}d_{jt}[E(\varepsilon_{jt}|d_t) - E(\varepsilon_{jt}|d_{jt} = 1)]\quad (9)$$

Second, we take the expectation with respect to d_t , yielding

$$E(\xi_{it}|x_t) = \sum_{\substack{j=1 \\ j \neq i}}^m \omega_{ijt}\Pr(d_{it}d_{jt} = 1)[E(\varepsilon_{jt}|d_{it}d_{jt} = 1) - E(\varepsilon_{jt}|d_{jt} = 1)].\quad (10)$$

From (10) it becomes obvious that the estimator $\widehat{\beta}$ is biased and inconsistent unless either (i) $E(\varepsilon_{jt}|d_{it}d_{jt} = 1)$ equals $E(\varepsilon_{jt}|d_{jt} = 1)$ for any pair $i \neq j$ and any t ; that is $E(\varepsilon_{it}|d_t)$ exclusively depends on d_{it} , yet does not depend on any d_{jt} , $j \neq i$. This requires that Σ_{vv} as well as $\Sigma_{\varepsilon v}$ are diagonal matrices. The estimator $\widehat{\beta}$ does also not suffer from inconsistency if (ii) $\omega_{ijt} = 0$ holds for all $i \neq j$ and t . Condition (ii) implies that equation-by-equation Heckman is consistent, since cross-equation correlations are not taken into account. Yet, in contrast, any system estimator that involves non-zero

weights ω_{ijt} is inconsistent, unless Σ_{vv} as well as $\Sigma_{\varepsilon v}$ are diagonal matrices. Clearly, the inconsistency of $\hat{\beta}$ originates from conditioning on d_{it} equation-by-equation.

3 A consistent generalized Heckman estimator

The above discussion clearly suggests, how to construct a consistent system-estimator as generalization to the original Heckman-approach. From (9) follows that if $\tilde{\varepsilon}_{it}$ were defined as $\varepsilon_{it} - E(\varepsilon_{it}|d_t)$ rather than $\varepsilon_{it} - E(\varepsilon_{it}|d_{it} = 1)$, the condition $E(\xi_{it}|x_t, d_t) = 0$ and subsequently $E(\xi_{it}|x_t) = 0$ would be satisfied for any weighting matrix Ω , rendering the entire class of estimators consistent. Uniformly conditioning on d_t , i.e. conditioning on the entire selection pattern, in all equations rather than conditioning on d_{it} equation-by-equation and, correspondingly, including $E(\varepsilon_{it}|d_t)$ rather than the inverse Mills ratio as correction-term would lead to errors defined as $\varepsilon_{it} - E(\varepsilon_{it}|d_t)$. That is, the regression must be based on the conditional mean $E(y_{it}|x_{it}, d_t)$ rather than $E(y_{it}|x_{it}, d_{it})$.

In order to implement this estimator, an expression for $E(\varepsilon_{it}|d_t)$ is required. It is easily shown that

$$E(\varepsilon_t|d_t) = E(\varepsilon_t) + \Sigma_{\varepsilon v}(\Sigma_{vv})^{-1}[E(v_t|d_t) - E(v_t)] \quad (11)$$

holds. Since the unconditional expectations of ε_t and v_t equal zero, the expression reduces to a linear-combination of truncated first moments $E(v_t|d_t)$ from the multivariate normal distribution. Therefore, in each regression equation m truncated means

from the multivariate normal distribution have to be included to correct for sample-selection bias. Results for these truncated means are provided by Tallis (1961) as well as for the special case $m = 2$ – albeit in more detail – by Shah & Parikh (1964). Including these expressions and rearranging terms leads to the system of regression equations

$$y_{it} = d_{it}x'_{it}\beta_i + d_{it} \sum_{j=1}^m \delta_{ij} \psi_{jt} \phi(z'_{jt}\alpha_j) \frac{\Phi^{m-1}(\tilde{A}_{jt}, \tilde{R}_{jt})}{\Phi^m(\bullet)} + d_{it}\tilde{\varepsilon}_{it}, \quad i = 1, \dots, m. \quad (12)$$

As in the original Heckman-model, the coefficients δ_{ij} attached to the correction-terms $\psi_{jt}\phi(z'_{jt}\alpha_j) \frac{\Phi^{m-1}(\tilde{A}_{jt}, \tilde{R}_{jt})}{\Phi^m(\bullet)}$ are subject to estimation. Here, ϕ denotes the probability density function of the univariate standard normal distribution, while Φ^m denotes the cumulative density function of the m -variate standard normal distribution. ψ_{jt} is defined as $2d_{jt} - 1$ and distinguishes truncation from either below or above. \tilde{A}_{jt} represents a vector which consists of $m - 1$ elements $\frac{\psi_{lt}(z'_{lt}\alpha_l - \sigma_{lj}^{vv} z'_{jt}\alpha_j)}{(1 - (\sigma_{lj}^{vv})^2)^{1/2}}$; $l = 1 \dots m, l \neq j$. Correspondingly, \tilde{R}_{jt} is defined as $\Psi_{jt}R_{jt}\Psi_{jt}$, where R_{jt} denotes the partial conditional correlation-matrix $\text{Cor}(v_t|v_{jt})$ and Ψ_{jt} denotes a diagonal-matrix with diagonal elements $\psi_{lt}, l \neq j$. Finally, $\Phi^m(\bullet)$ denotes the joint probability of the observed pattern d_t . Note that the regression equations are still weighted by d_{it} .⁷

In applied work α and Σ_{vv} are likely to be unknown. In order to calculate the auxiliary regressors $\psi_{jt}\phi(z'_{jt}\alpha_j) \frac{\Phi^{m-1}(\tilde{A}_{jt}, \tilde{R}_{jt})}{\Phi^m(\bullet)}$, one has to replace the true parameters with estimates obtained from first-step multivariate probit estimation. In the special

⁷Since $E(y_{it}|x_{it}, d_t, d_{it} = 0) = 0$ holds, the i th equation of the t th observation still receives zero weight if y_{it} equals zero because of censoring.

case $m = 2$ the regression equations are equivalent to the one used by Poirier (1980), except for the fact that Poirier (1980) conditions on $d_{1t}d_{2t} = 1$ rather than d_{1t} and d_{2t} , i.e. ψ_{jt} equals one for all j and t .⁸ For $m = 2$, $\delta_{ij} = \sigma_{ij}^{\varepsilon v}$ holds for the auxiliary regression coefficients.

One may estimate the system (12) equation-by-equation using *OLS*. Yet, the simple equation-by-equation Heckman-estimator is consistent as well in this case. So, conditioning on d_t makes sense only in the context of simultaneous estimation. As a simple variant, one can construct such a system-estimator in the standard *SUR* fashion. However, this ignores the heteroscedasticity of the individual conditional error-variances. In order to be able to construct a proper *GLS* estimator, expressions for $\text{Var}(\varepsilon_t|d_t)$ are required from which one can calculate an appropriate weighting matrix Ω . Through the use of the normality assumption and the decomposition rule for variances in a joint distribution such an expression can easily be derived as

$$\text{Var}(\varepsilon_t|d_t) = \Sigma_{\varepsilon\varepsilon} - \Sigma_{\varepsilon v}(\Sigma_{vv})^{-1}\Sigma'_{\varepsilon v} + \Sigma_{\varepsilon v}(\Sigma_{vv})^{-1}\text{Var}(v_t|d_t)(\Sigma_{vv})^{-1}\Sigma'_{\varepsilon v}. \quad (13)$$

Obviously, any element of $\text{Var}(\varepsilon_t|d_t)$ is a linear function of all elements of the truncated m -variate normal variance-covariance matrix $\text{Var}(v_t|d_t)$. Therefore, estimates for the elements of $\text{Var}(\varepsilon_t|d_t)$ can be obtained as fitted values from regressing squared residuals and residual cross-products – which, in turn, have been obtained from initial *OLS* regressions – on a constant and on estimates for all elements of $\text{Var}(v_t|d_t)$.⁹ Results

⁸See Vella (1997) for other related models.

⁹Because of $\text{var}(d_{it}\varepsilon_{it}|d_{it} = 0) = 0$, the variance-covariance matrix $\text{Var}(d_{1t}\varepsilon_{1t} \dots d_{mt}\varepsilon_{mt}|d_t)$ that

for the latter ones are provided by Tallis (1961). Therefore, with estimates for α and Σ_{vv} in hand, one can calculate these auxiliary regressors.

4 Monte-Carlo analysis

In addition to the theoretical analysis we carry out Monte-Carlo simulations. On the one hand, we want to illustrate the theoretical results derived in Section 2. Test results on the joint unbiasedness of the second-step coefficients are provided for this purpose.¹⁰

On the other, we also want to address the estimators' performance beyond the issue of consistency. Therefore, we present estimates for the *CP*-conditional mean square error prediction criterion

$$\text{CP}(\hat{\beta}) = \text{E} \left[\frac{1}{T} \sum_{t=1}^T \sum_{i=1}^m (\beta_i - \hat{\beta}_i)' x_{it} x_{it}' (\beta_i - \hat{\beta}_i) \middle| X \right], \quad (14)$$

cf. Judge et al. (1980). $\text{CP}(\hat{\beta})$ measures the mean squared deviation of the estimated conditional mean from its true counterpart $\text{E}(y_{it}^* | x_{it})$ and, therefore, translates an estimator's *MSE*-matrix to a scalar performance measure that takes into account its variance as well as a potential bias.

is effectively required for the construction of the *GLS* estimator in general is short-ranked and cannot ordinarily be inverted in order to obtain individual weighting-matrices Ω_t . Yet, using a generalized Moore-Penrose inverse is appropriate for this purpose.

¹⁰Both the tables of raw coefficients' estimates as well as the *LIMDEP* command file used for carrying out the *MC*-simulations are available from the author upon request.

Unknown values for α and Σ_{vv} rather than known ones appear to be the relevant case from the viewpoint of applied econometrics. In our Monte-Carlo simulations, therefore, these parameters are estimated by first-step probit models. We consider six different estimators. In particular, conditioning on either d_{it} or d_t is combined with *OLS*, *SUR* and, finally *GLS* estimation. Conditioning on d_{it} combined with *OLS* or *SUR* allows for estimating the first step using univariate probit models. All other estimators require simultaneous estimation of all vectors α_i along with Σ_{vv} .

4.1 The experimental setup

The design of the Monte-Carlo experiment is equivalent to the one used by Tauchmann (2005).¹¹ We consider the case $m = 2$.¹² The sample size is 4000. The size of the Monte Carlo experiment is 1000 iterations. The vectors of exogenous variables each consist of three elements:

$$z_{it} = [1 \ z_{1,it} \ z_{2,it}]', \quad x_{it} = [1 \ x_{1,it} \ x_{2,it}]', \quad i = 1, 2.$$

Here $z_{1,1t}$, $z_{2,1t}$, $z_{1,2t}$, and $x_{2,1t}$ are independently drawn from the standard normal distribution, while $z_{2,2t} = z_{2,1t}$, $x_{1,1t} = z_{1,1t}$, $x_{1,2t} = z_{1,2t}$ and $x_{2,2t} = x_{2,1t}$. These variables are drawn only once and then kept fixed. For the coefficient vectors $\beta_i = [1 \ 1 \ 1]'$,

¹¹In contrast to the analysis presented here, Tauchmann (2005) imposes restrictions on the coefficients' estimates $\hat{\beta}_i$. This does not allow for directly comparing estimated *CP*-measures.

¹²For $m \geq 3$, simulated *ML* were required for estimation the first-step multivariate probit models. This would increase computing time for the Monte-Carlo experiments enormously.

$i = 1, 2$ holds.¹³

The value $\sqrt{0.5}$ is assigned to all coefficients α attached to $z_{1,it}$ and $z_{2,it}$. In order to allow for different unconditional censoring probabilities $\Pr(d_{it}^* \leq 0)$, the constants $\alpha_{0,i}$ are varied. We run two simulations with unconditional censoring probabilities that are uniform across equations, in particular 0.25 and 0.5, which corresponds to constants 0.9539 and 0, respectively. Another simulation is carried out for mixed unconditional censoring probabilities, i.e. 0.25 for equation one and 0.75 for equation two, which corresponds to constants 0.9539 and -0.9539 , respectively. The error-covariance structure is specified as

$$\Sigma_{\varepsilon\varepsilon} = \begin{bmatrix} 1.5 & \\ -1 & 2 \end{bmatrix}, \quad \Sigma_{vv} = \begin{bmatrix} 1 & \\ -0.5 & 1 \end{bmatrix}, \quad \Sigma_{\varepsilon v} = \begin{bmatrix} 0.75 & -0.25 \\ -0.25 & 0.75 \end{bmatrix}.$$

As an alternative specification, the value zero is assigned to all off-diagonal elements of Σ_{vv} and $\Sigma_{\varepsilon v}$ everything else remaining unchanged, i.e.

$$\Sigma_{\varepsilon\varepsilon} = \begin{bmatrix} 1.5 & \\ -1 & 2 \end{bmatrix}, \quad \Sigma_{vv} = \begin{bmatrix} 1 & \\ 0 & 1 \end{bmatrix}, \quad \Sigma_{\varepsilon v} = \begin{bmatrix} 0.75 & 0 \\ 0 & 0.75 \end{bmatrix}.$$

This defines the four-variate $N(0, \Sigma)$ distribution, from where the random components are drawn separately for each model. After drawing the error vector, the dependent variables are calculated as defined by model (1) through (4). Subsequently, the generated data serves as input to the estimators.

¹³We do not vary these parameters, since – in contrast to the estimator proposed by Shonkwiler & Yen (1999) – the performance of generalized Heckman estimators does not depend on the true value of β , c.f. Tauchmann (2005).

4.2 Simulation results

Results for Wald-tests on the unbiasedness of the six estimators are displayed in Table 1. These simulation results are consistent with the theoretical ones, obtained in Section 2. If Σ_{vv} and $\Sigma_{\varepsilon v}$ are dense matrices, unbiasedness is clearly rejected for those estimators that condition on d_{it} equation-by-equation and use *SUR* or *GLS*. In contrast, the classical Heckman estimator employed equation-by-equation does not display a significant bias. The estimators that condition on the entire selection pattern do not display a significant bias either. If, instead, Σ_{vv} and $\Sigma_{\varepsilon v}$ are diagonal-matrices, neither of the estimators display a bias that is significant at the 0.05-level. Therefore, the Monte-Carlo simulation confirms that system-estimators that condition on d_t are consistent, while system-estimators that condition on d_{it} equation-by-equation are biased, unless certain restrictions apply to the true error-covariance matrix.

< insert Table 1 about here >

In order to analyze the estimators' performance beyond the issue of unbiasedness, estimates for the *CP*-conditional mean square error prediction criterion are displayed in Table 2. Comparing the *SUR* estimator that conditions on d_t with its counterpart that conditions on d_{it} yields the following plausible result: If the true covariance-matrix is dense, the consistent estimator that conditions on d_t yields smaller *CP*-measures than the inconsistent one that conditions on d_{it} . If Σ_{vv} and $\Sigma_{\varepsilon v}$ are diagonal-matrices – i.e. both estimators are consistent – the more parsimoniously parameterized one that conditions on d_{it} performs better except for one simulation. Yet, the latter differences

1
2
3
4 in estimated CP -measures are statistically insignificant at the 0.05-level.
5
6

7 The comparison of OLS estimators that either condition on d_t or d_{it} yields similar
8 results. If the error-covariance matrix is dense, the first estimator seems to perform
9 better, though both are consistent. If, instead, Σ_{vv} and $\Sigma_{\varepsilon v}$ are diagonal-matrices the
10 latter displays smaller CP -measures. However, these differences never are statistically
11 significant, except for one simulation.
12
13
14
15
16
17

18
19
20
21 < insert Table 2 about here >
22
23

24 Finally, we examine the performance of GLS estimators. Here, we observe sub-
25 stantial deviations in estimated CP -measures. While GLS conditional on d_t yields the
26 smallest mean square prediction error among all considered estimators in any simu-
27 lation, GLS conditional on d_{it} , except for two simulations, displays the largest one.
28 Moreover, the deviations in CP -measures between both GLS estimators always are
29 significant. In fact, if the error covariance-matrix is dense, GLS conditional on d_t sig-
30 nificantly outperforms any other estimator in any simulation. As the only exception
31 to this result, in some cases SUR conditional on d_t displays CP -measures which are
32 not significantly larger.
33
34
35
36
37
38
39
40
41
42
43
44
45

46 Our key simulation result – that GLS conditional on d_t displays the best perfor-
47 mance in terms of the mean square prediction error – fits theory. Among the considered
48 estimators, GLS conditional on d_t is the only one that not only is consistent, but also
49 properly accounts for cross-equation correlation and heteroscedasticity.
50
51
52
53
54
55
56
57
58
59
60

5 Conclusions

This analysis of estimation procedures for the multivariate sample-selection model shows that multivariate generalizations to the classical Heckman (1976 and 1979) two-step approach that account for cross-equation correlation and use the inverse Mills ratio as a correction-term are consistent only if certain restrictions apply to the true error-covariance structure. However, generalizations to the classical Heckman two-step estimator that condition on the entire selection pattern rather than the selection of particular single equations – and, therefore, use generalized correction-terms – are shown to be generally consistent. Moreover, these estimators display a smaller mean square prediction error. These new estimators are computationally more demanding since they generally require simultaneous estimation of a multivariate probit model. Nowadays, however, hard-coded procedures for this estimation problem are provided by econometric software packages, rendering computational complexity a minor obstacle to the practical application of the suggested estimation procedure.

Finally, we discuss how our results fit into the general debate on which estimator is the best choice for estimating the multivariate sample selection model. If efficiency is the major concern and numerical complexity and computing time do not matter, then two-step approaches – including those suggested in this analysis – are generally to be avoided, and full information maximum likelihood as proposed by Yen (2005) is the best choice. If, in contrast, computational simplicity and consistency is the major concern, then equation-by-equation Heckman appears to be the best choice. If a small

1
2
3
4 mean square error and computational simplicity are a researcher's main criteria, while
5
6 consistency is of secondary relevance, one might even argue in favor of the inconsistent
7
8 *SUR* estimator that conditions equation-by-equation on the outcome of the upstream
9
10 choice problem. Finally, if both consistency and a small mean square error are desired,
11
12 and the computational burden of full information maximum likelihood is to be avoided,
13
14 then the *GLS* estimator that conditions on the entire selection pattern appears to be
15
16 the best choice.
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

References

- 1
2
3
4
5
6
7
8 **Chen Z. and S.T. Yen** (2005): On bias correction in the multivariate sample-
9 selection model, *Applied Economics* 37, 2459-2468.
10
11
12
13
14 **Gao X.M. E.J. Wailes and G.L. Cramer** (1995): A Microeconomic Model
15 Analysis of US Consumer Demand for Alcoholic Beverages, *Applied Economics*
16 27, 59-69.
17
18
19
20
21
22 **Goodwin B.K. M.L. Vandever and J.L. Deal** (2004): An Empirical Analy-
23 sis of Acreage Effects of Participation in the Federal Crop Insurance Program,
24 *American Journal of Agricultural Economics* 86, 1058-1077.
25
26
27
28
29
30
31 **Heckman J.J.** (1976): The Common Structure of Statistical Models of Truncation,
32 Sample Selection and Limited Dependent Variables and a Simple Estimator for
33 such Models, *Annals of Economics and Social Measurement* 5, 475-492.
34
35
36
37
38
39 **Heckman J.J.** (1979): Sample Selection Bias as a Specification Error, *Econometrica*
40 47, 153-161.
41
42
43
44
45 **Heien D. and C. Durham** (1991): A Test of the Habit Formation Hypothesis
46 using Household Data, *Review of Economics and Statistics* 73, 189-199.
47
48
49
50
51 **Heien D. and C.R. Wessells** (1990): Demand Systems Estimation with Microdata:
52 A Censored Regression Approach, *Journal of Business & Economic Statistics* 8,
53 365-371.
54
55
56
57
58
59
60

1
2
3
4 **Judge G.G. et al.** (1980): The Theory and Practice of Econometrics, 2nd ed., John
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Wiley, New York.

Nayga R.M. B.J. Tepper and L. Rosenzweig (1999): Assessing the Importance
of Health and Nutrition related Factors on Food Demand: a Variable Preference
Investigation, *Applied Economics* 31, 1541-1549.

Poirier D.J. (1980): Partial Observability in Bivariate Probit Models, *Journal of
Econometrics* 12, 209-217.

Shah S.M. and N.T. Parikh (1964): Moments of Singly and Doubly Truncated
Standard Bivariate Normal Distribution, *Vidya* 7, 51-91.

Shonkwiler J.S. and S.T. Yen (1999): Two-Step Estimation of a Censored System
of Equations, *American Journal of Agricultural Economics* 81, 972-982.

Su S.-J. and S.T. Yen (2000): A Censored System of Cigarette and Alcohol Con-
sumption, *Applied Economics* 32, 729-737.

Tallis G.M. (1961): The Moment Generating Function of the Truncated Multi-
Normal Distribution, *Journal of the Royal Statistical Society* (Series B) 23, 223-
229.

Tauchmann H. (2005): Efficiency of two-step Estimators for Censored Systems of
Equations: Shonkwiler and Yen reconsidered, *Applied Economics* 37, 367-374.

1
2
3
4 **Vella F.** (1997): Estimating Models with Sample Selection Bias: A Survey, *Journal*
5
6 *of Human Resources* 33, 127-169.
7
8

9
10 **Vermeulen F.** (2001): A Note on Heckman-type Corrections in Models for Zero
11
12 Expenditures, *Applied Economics* 33, 1089-1092.
13
14

15
16 **Yen S.T. K. Kah and S.-J. Su** (2002): Household Demand for Fat and Oils:
17
18 Two-step Estimation of a Censored Demand System, *Applied Economics* 34,
19
20 1799-1806.
21
22

23
24 **Yen S.T.** (2005): A Multivariate Sample-Selection Model: Estimating Cigarette
25
26 and Alcohol demand with Zero Observations, *American Journal of Agricultural*
27
28 *Economics* 87, 453-466.
29
30

31
32 **Zellner A.** (1963): An Efficient Method of Estimating Seemingly Unrelated Regres-
33
34 sion Equations: Some Exact Finite Sample Results, *Journal of the American*
35
36 *Statistical Association* 58, 977-992.
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Table 1: Tests on joint unbiasedness of regression coefficients

	<i>OLS</i>	<i>SUR</i>	<i>GLS</i>	<i>OLS</i>	<i>SUR</i>	<i>GLS</i>
	conditional on d_{it}			conditional on d_t		
dense error variance-covariance matrix						
censoring prob. 0.25	0.800	0.000	0.000	0.449	0.484	0.155
censoring prob. 0.5	0.545	0.000	0.000	0.964	0.070	0.642
censoring prob. 0.25 and 0.75	0.446	0.000	0.000	0.117	0.929	0.259
Σ_{vv} and $\Sigma_{\varepsilon v}$ with zero off-diagonal elements						
censoring prob. 0.25	0.320	0.415	0.052	0.805	0.208	0.082
censoring prob. 0.5	0.595	0.659	0.610	0.900	0.760	0.807
censoring prob. 0.25 and 0.75	0.832	0.620	0.604	0.963	0.634	0.215

Note: P-values for Wald-tests reported.

Table 2: Estimated conditional mean square prediction errors

	<i>OLS</i>	<i>SUR</i>	<i>GLS</i>	<i>OLS</i>	<i>SUR</i>	<i>GLS</i>
	conditional on d_{it}			conditional on d_t		
dense error variance-covariance matrix						
censoring prob. 0.25	6.499 (0.154)	6.038 (0.149)	6.455 (0.169)	5.826 (0.130)	5.468 (0.137)	5.357 (0.140)
censoring prob. 0.5	12.810 (0.342)	13.444 (0.377)	15.816 (0.483)	12.275 (0.309)	11.776 (0.335)	11.178 (0.330)
cens. prob. 0.25 & 0.75	23.747 (0.806)	23.129 (0.755)	35.492 (1.391)	23.095 (0.784)	21.877 (0.762)	19.387 (0.786)
Σ_{vv} and $\Sigma_{\varepsilon v}$ with zero off-diagonal elements						
censoring prob. 0.25	6.420 (0.156)	5.520 (0.135)	5.872 (0.212)	6.304 (0.147)	5.452 (0.138)	5.269 (0.140)
censoring prob. 0.5	13.451 (0.386)	11.776 (0.346)	15.880 (1.601)	13.533 (0.364)	11.932 (0.324)	11.702 (0.352)
cens. prob. 0.25 & 0.75	23.219 (0.768)	20.627 (0.744)	28.690 (2.733)	24.156 (0.853)	21.091 (0.710)	17.748 (0.655)

Notes: Standard errors in parenthesis.

Displayed *CP*-measures are scaled by the factor 1000.

Estimated coefficients from Monte-Carlo simulations

Table 1: Estimated coefficients: censoring prob. 0.25, dense error variance-covariance matrix

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	1.0002	0.0389	0.9881	0.0373	0.9859	0.0373	0.9994	0.0363	1.0009	0.0361	1.0014	0.0363
$\beta_{1,1}$	1	1.0000	0.0285	1.0041	0.0253	1.0041	0.0254	0.9989	0.0280	1.0000	0.0248	0.9991	0.0249
$\beta_{2,1}$	1	0.9993	0.0209	1.0007	0.0203	1.0005	0.0202	0.9997	0.0206	0.9993	0.0202	0.9986	0.0205
$\sigma_{11}^{\varepsilon}$	0.75	0.7488	0.0900	0.7459	0.0873	0.7464	0.0868	0.7489	0.0848	0.7464	0.0813	0.7459	0.0830
$\sigma_{12}^{\varepsilon}$	-0.25	-	-	-	-	-	-	-0.2505	0.0369	-0.2496	0.0375	-0.2499	0.0362
$\beta_{0,2}$	1	1.0015	0.0458	0.9850	0.0427	0.9756	0.0424	1.0020	0.0413	1.0001	0.0409	0.9993	0.0419
$\beta_{1,2}$	1	1.0003	0.0325	1.0071	0.0285	1.0092	0.0283	1.0004	0.0317	1.0016	0.0303	1.0005	0.0290
$\beta_{2,2}$	1	1.0002	0.0246	0.9993	0.0238	0.9998	0.0234	1.0011	0.0250	1.0003	0.0240	0.9992	0.0241
$\sigma_{22}^{\varepsilon}$	0.75	0.7488	0.1079	0.7475	0.0953	0.7548	0.0979	0.7466	0.0979	0.7513	0.0930	0.7509	0.0956
$\sigma_{21}^{\varepsilon}$	-0.25	-	-	-	-	-	-	-0.2495	0.0433	-0.2512	0.0426	-0.2490	0.0430

Table 2: Estimated coefficients: censoring prob. 0.5, dense error variance-covariance matrix

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	0.9963	0.0596	0.9886	0.0586	0.9670	0.0641	0.9993	0.0587	1.0014	0.0604	0.9994	0.0586
$\beta_{1,1}$	1	1.0016	0.0351	1.0036	0.0339	1.0079	0.0346	1.0006	0.0361	0.9992	0.0334	1.0008	0.0327
$\beta_{2,1}$	1	1.0006	0.0250	1.0001	0.0249	1.0000	0.0234	0.9995	0.0256	1.0013	0.0245	0.9999	0.0239
$\sigma_{11}^{\varepsilon v}$	0.75	0.7543	0.0783	0.7579	0.0763	0.7793	0.0846	0.7507	0.0759	0.7455	0.0765	0.7507	0.0736
$\sigma_{12}^{\varepsilon v}$	-0.25	-	-	-	-	-	-	-0.2492	0.0411	-0.2466	0.0410	-0.2513	0.0402
$\beta_{0,2}$	1	1.0031	0.0699	0.9803	0.0734	0.9628	0.0707	1.0003	0.0681	0.9988	0.0650	1.0037	0.0650
$\beta_{1,2}$	1	0.9984	0.0413	1.0079	0.0398	1.0087	0.0418	1.0008	0.0380	1.0005	0.0382	0.9987	0.0369
$\beta_{2,2}$	1	1.0008	0.0295	0.9999	0.0278	0.9990	0.0269	0.9998	0.0294	0.9989	0.0282	0.9998	0.0278
$\sigma_{22}^{\varepsilon v}$	0.75	0.7472	0.0911	0.7660	0.0977	0.7824	0.0917	0.7490	0.0879	0.7511	0.0867	0.7482	0.0829
$\sigma_{21}^{\varepsilon v}$	-0.25	-	-	-	-	-	-	-0.2510	0.0476	-0.2476	0.0477	-0.2489	0.0449

Table 3: Estimated coefficients: censoring prob. 0.25 and 0.75, dense error variance-covariance matrix

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	0.9984	0.0396	0.9975	0.0386	0.9963	0.0385	1.0001	0.0373	1.0005	0.0363	1.0005	0.0378
$\beta_{1,1}$	1	1.0011	0.0283	1.0010	0.0278	0.9964	0.0281	1.0009	0.0272	0.9995	0.0274	0.9990	0.0278
$\beta_{2,1}$	1	0.9991	0.0207	1.0000	0.0205	0.9999	0.0213	1.0004	0.0211	1.0008	0.0215	0.9993	0.0209
σ_{11}^{EV}	0.75	0.7550	0.0937	0.7537	0.0915	0.7592	0.0948	0.7489	0.0850	0.7490	0.0848	0.7498	0.0852
σ_{12}^{EV}	-0.25	-	-	-	-	-	-	-0.2485	0.0354	-0.2484	0.0351	-0.2505	0.0351
$\beta_{0,2}$	1	1.0060	0.1275	0.9725	0.1240	0.8927	0.1262	1.0057	0.1259	0.9992	0.1235	0.9983	0.1171
$\beta_{1,2}$	1	0.9979	0.0566	1.0090	0.0542	1.0311	0.0560	1.0014	0.0572	1.0009	0.0527	1.0022	0.0529
$\beta_{2,2}$	1	1.0005	0.0410	0.9998	0.0391	0.9983	0.0394	0.9992	0.0401	0.9996	0.0391	0.9984	0.0374
σ_{22}^{EV}	0.75	0.7440	0.1058	0.7564	0.1014	0.7775	0.1113	0.7444	0.1056	0.7495	0.1011	0.7533	0.0981
σ_{21}^{EV}	-0.25	-	-	-	-	-	-	-0.2468	0.0745	-0.2493	0.0720	-0.2496	0.075

Table 4: Estimated coefficients: censoring prob. 0.25, diagonal-matrices Σ_{uv} and $\Sigma_{\varepsilon v}$

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	0.9985	0.0626	1.0025	0.0602	0.9978	0.0704	1.0005	0.0608	0.9997	0.0585	0.9978	0.0592
$\beta_{1,1}$	1	1.0012	0.0355	0.9984	0.0326	1.0014	0.0395	0.9995	0.0356	1.0009	0.0323	1.0008	0.0327
$\beta_{2,1}$	1	1.0010	0.0249	1.0002	0.0231	0.9997	0.0241	1.0005	0.0258	1.0000	0.0242	0.9988	0.0238
$\sigma_{11}^{\varepsilon v}$	0.75	0.7528	0.0787	0.7464	0.0774	0.7543	0.1000	0.7511	0.0797	0.7494	0.0728	0.7522	0.0742
$\sigma_{12}^{\varepsilon v}$	0	-	-	-	-	-	-	0.0012	0.0372	0.0007	0.0369	0.0000	0.0367
$\beta_{0,2}$	1	1.0015	0.0721	1.0003	0.0668	0.9958	0.0771	0.9974	0.0730	1.0004	0.0693	1.0029	0.0685
$\beta_{1,2}$	1	0.9991	0.0410	0.9993	0.0373	1.0025	0.0460	1.0007	0.0422	0.9990	0.0379	0.9997	0.0368
$\beta_{2,2}$	1	0.9997	0.0295	0.9992	0.0277	0.9995	0.0288	1.0001	0.0297	0.9989	0.0271	1.0008	0.0283
$\sigma_{22}^{\varepsilon v}$	0.75	0.7498	0.0946	0.7487	0.0854	0.7566	0.1076	0.7524	0.0959	0.7505	0.0891	0.7474	0.0871
$\sigma_{21}^{\varepsilon v}$	0	-	-	-	-	-	-	-0.0009	0.0427	0.0009	0.0429	0.0010	0.0447

Table 5: Estimated coefficients: censoring prob. 0.5, diagonal-matrices Σ_{vv} and $\Sigma_{\varepsilon v}$

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	0.9985	0.0626	1.0025	0.0602	0.9978	0.0704	1.0005	0.0608	0.9997	0.0585	0.9978	0.0592
$\beta_{1,1}$	1	1.0012	0.0355	0.9984	0.0326	1.0014	0.0395	0.9995	0.0356	1.0009	0.0323	1.0008	0.0327
$\beta_{2,1}$	1	1.0010	0.0249	1.0002	0.0231	0.9997	0.0241	1.0005	0.0258	1.0000	0.0242	0.9988	0.0238
$\sigma_{11}^{\varepsilon v}$	0.75	0.7528	0.0787	0.7464	0.0774	0.7543	0.1000	0.7511	0.0797	0.7494	0.0728	0.7522	0.0742
$\sigma_{12}^{\varepsilon v}$	0	-	-	-	-	-	-	0.0012	0.0372	0.0007	0.0369	0.0000	0.0367
$\beta_{0,2}$	1	1.0015	0.0721	1.0003	0.0668	0.9958	0.0771	0.9974	0.0730	1.0004	0.0693	1.0029	0.0685
$\beta_{1,2}$	1	0.9991	0.0410	0.9993	0.0373	1.0025	0.0460	1.0007	0.0422	0.9990	0.0379	0.9997	0.0368
$\beta_{2,2}$	1	0.9997	0.0295	0.9992	0.0277	0.9995	0.0288	1.0001	0.0297	0.9989	0.0271	1.0008	0.0283
$\sigma_{22}^{\varepsilon v}$	0.75	0.7498	0.0946	0.7487	0.0854	0.7566	0.1076	0.7524	0.0959	0.7505	0.0891	0.7474	0.0871
$\sigma_{21}^{\varepsilon v}$	0	-	-	-	-	-	-	-0.0009	0.0427	0.0009	0.0429	0.0010	0.0447

Table 6: Estimated coefficients: censoring prob. 0.25 and 0.75, diagonal-matrices Σ_{vv} and $\Sigma_{\varepsilon v}$

true value	conditional on d_{it}						conditional on d_t						
	OLS		SUR		GLS		OLS		SUR		GLS		
	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	mean	st. dev.	
$\beta_{0,1}$	1	0.9998	0.0395	0.9984	0.0385	1.0004	0.0443	1.0009	0.0383	1.0012	0.0373	1.0025	0.0385
$\beta_{1,1}$	1	1.0006	0.0281	0.9999	0.0272	1.0001	0.0338	1.0003	0.0281	0.9999	0.0274	0.9994	0.0269
$\beta_{2,1}$	1	1.0000	0.0212	0.9996	0.0206	1.0006	0.0215	0.9999	0.0211	1.0005	0.0206	0.9990	0.0208
$\sigma_{11}^{\varepsilon v}$	0.75	0.7502	0.0900	0.7524	0.0876	0.7502	0.1129	0.7467	0.0880	0.7480	0.0865	0.7454	0.0892
$\sigma_{12}^{\varepsilon v}$	0	-	-	-	-	-	-	-0.0001	0.0339	0.0001	0.0336	-0.0003	0.0335
$\beta_{0,2}$	1	0.9991	0.1267	0.9998	0.1178	0.9994	0.1393	1.0000	0.1288	0.9971	0.1205	1.0037	0.1114
$\beta_{1,2}$	1	1.0013	0.0557	0.9984	0.0530	0.9988	0.0652	1.0001	0.0586	0.9997	0.0531	0.9999	0.0491
$\beta_{2,2}$	1	0.9992	0.0386	1.0000	0.0386	0.9993	0.0419	1.0000	0.0411	0.9989	0.0374	1.0004	0.0354
$\sigma_{22}^{\varepsilon v}$	0.75	0.7523	0.1056	0.7506	0.0958	0.7531	0.1234	0.7502	0.1077	0.7507	0.0982	0.7454	0.0911
$\sigma_{21}^{\varepsilon v}$	0	-	-	-	-	-	-	0.0010	0.0743	0.0034	0.0717	0.0005	0.0756

Seeds for the Random Numbers Generator

Table 7: Seeds for the Random Numbers Generator

<i>MC-Simulation</i>	Random Seed
dense error variance-covariance matrix	
censoring prob. 0.25	61925187
censoring prob. 0.5	92306363
cens. prob. 0.25 & 0.75	99416675
Σ_{vv} and $\Sigma_{\varepsilon v}$ with zero off-diagonal elements	
censoring prob. 0.25	64300457
censoring prob. 0.5	61668177
cens. prob. 0.25 & 0.75	32974293

1 Consig_CompEconsworld_Klein_2010G.Lim
2 Friday, 17 March 2006

Page 1
14:12:00 AM

3 ?? LIMDEP COMMAND-FILE USED FOR MC-SIMULATIONS

4 ?? GENERAL PARAMETERS

5 ?? Sample Size

6 CALC ; NN= 4000 \$
7 Sample ; 1-NN \$

8 ?? Parameter Values

9 ?? Error-Covariance-Matrix "MUU"

10 MATRIX ; MU11= [1/-0.5,1]
11 ; MU22= [1.5/-1,2]
12 ; MU12= [0.75,-0.25/-0.25,0.75] \$

13 MATRIX ; MU21= MU12'
14 ; MUU=[MU11,MU12/MU21,MU22] \$

15 CALC ; cor= MUU(1,2) \$

16 ?? Cholesky-Factorisation

17 MATRIX ; LMU= Chol(MUU)
18 ; LMU1= Part(LMU,1,1,1,4)
19 ; LMU2= Part(LMU,2,2,1,4)
20 ; LMU3= Part(LMU,3,3,1,4)
21 ; LMU4= Part(LMU,4,4,1,4) \$

22 ?? Censoring Probability

23 CALC ; CENSPRO1= 0.25
24 ; CENSPRO2= 0.25 \$

25 ?? Coefficients: Discrete Model

26 Calc ; alf11 = (0.5)^0.5
27 ; alf21= (0.5)^0.5
28 ; list ; ALF01= -(1+ alf11^2+ alf21^2)^0.5*Ntb(CENSPRO1)
29 ; list ; Alf02= -(1+ alf11^2+ alf21^2)^0.5*Ntb(CENSPRO2) \$

30 ?? Coefficient Vectors

31 MATRIX ; ALF1= [alf01/alf11/alf21]
32 ; ALF2= [alf02/alf11/alf21]
33 ; BET1= [1/1/1]
34 ; BET2= [1/1/1] \$

35 ?? Number of Loops in MC Experiment

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 1
04/12/2010 10:00:00 AM

CALC ; M= 1000 \$

?? No Trace-File !

NOTRACE \$

??

?? DATA SIMULATION

?? Exogenous Variables

?? Seed for Random Numbers Generator

CALC ; Ran(9876543) \$

CREATE ; Z1= Rnn(0,1) ; Z2= Rnn(0,1) ; X1= Z1 ; X2= Rnn(0,1) \$

CREATE ; Q1= Rnn(0,1) ; Q2= Z2 ; O1= Q1 ; O2= X2 \$

NAMELIST ; Z= ONE,Z1,Z2
; X= ONE,X1,X2
; Q= ONE,Q1,Q2
; O= ONE,O1,O2 \$

?? Additional Name-Lists

CREATE ; M1=0 ; M2=0
; R1=0 ; R2= 0 ; R3=0 ; R4=0 \$

NAMELI ; L1= X,M1
; L2= O,M2
; VA= R1,R2,R3,R4 \$

?? New Seed for Random Numbers Generator

CALC ; Ran(61925187) \$

??

?? COMPARISON of MODELS

??

C:\pnp\Comp\Gauss\work\heck_001006.lin
 Friday, 17 March 2006

Page 1
 04:12:00 AM

```

1  ?? EQUATING by EQUATION (OLS) HECKMAN-Model (Model #1)
2
3  ?? MONTE-CARLO EXPERIMENT
4
5  ?? Vectors for Intermediate Output
6
7  ? Renew Name-Lists
8
9  NAMELIST ; Z= ONE,Z1,Z2
10             ; X= ONE,X1,X2
11             ; Q= ONE,Q1,Q2
12             ; O= ONE,O1,O2
13             ; L1= X,M1
14             ; L2= O,M2
15             ; VA= R1,R2,R3,R4 $
16
17 MATRIX ; BCOL=  Init(M,14,0) $
18
19 PROC
20     DO FOR ; monte ; j= 1,M $
21         LABEL ; 99 $
22             ?? Drawing Random Error "ER"
23
24             ?? Drawing independent N(0,1) Random Numbers
25
26
27             CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
28                 ; R3= Rnn(0,1) ; R4= Rnn(0,1) $
29
30             NAME ; VA= R1,R2,R3,R4 $
31
32             ?? Transformation to N(0,MU) distributed Random Vector
33
34             CREA ; ER1= VA'LMU1
35                 ; ER2= VA'LMU2
36                 ; ER3= VA'LMU3
37                 ; ER4= VA'LMU4 $
38
39             ?? Discrete Endogenous Variables
40
41             CREA ; YD1= ALF1'Z + ER1
42                 ; if(YD1>0) YD1=1 ; (Else) YD1=0
43                 ; YD2= ALF2'Q + ER2
44                 ; if(YD2>0) YD2=1 ; (Else) YD2=0 $
45
46             ?? Continuous Endogenous Variables
47

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 4
04/12/2016 10:00 AM

```

CREA ; YS1=YD1*(BET1'X + ER3)
      ; YS2=YD2*(BET2'O + ER4) $

?? MODEL ESTIMATION

?? Univariate Probit Models

PROBIT ; lhs= YD1 ; rhs= Z
      ; start= alf1 $

MATRIX ; BP1= B $           ? Coefficient Vector
CREATE ; M1= N01(BP1'Z)/Phi(BP1'Z) ? Correction Term
      ; P1= Phi(BP1'Z) $

CALC   ; exi= opt_exit+ exitcode $

PROBIT ; lhs= YD2 ; rhs= Q
      ; start= alf2 $

MATRIX ; BP2= B $           ? Coefficient Vector
CREATE ; M2= N01(BP2'Q)/Phi(BP2'Q) ? Correction Term
      ; P2= Phi(BP2'Q) $

GO TO ; 99 ; opt_exit+ exitcode+ exi # 0 $ ? Convergence Control

?? Weighted Variables

CREATE ; X1Y1= X1*YD1 ; X2Y1= X2*YD1 ; M1Y1= M1*YD1
      ; X1Y2= O1*YD2 ; X2Y2= O2*YD2 ; M2Y2= M2*YD2 $

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-Estimation

NAME ; L1= YD1,X1Y1,X2Y1,M1Y1
      ; L2= YD2,X1Y2,X2Y2,M2Y2 $

MATR ; OL= L1'L1
      ; OR= Init(4,4,0)
      ; UL= Init(4,4,0)
      ; UR= L2'L2 $

MATR ; YO= L1'YS1
      ; YU= L2'YS2
      ; XW= [OL,OR/UL,UR]
      ; YW= [YO/YU] $

```

C:\msys5\opt\bin\win32\bin\win32\bin
 Friday, 17 March 2006

Page 3
 14:22:00 AM

```

1      MATR ; BR= <XW'XW>*XW'YW $
2
3      ? Accumulation of Results
4
5      MATR ; COEF= [BP1/BP2/BR]
6      ; BCOL(j,*)= COEF $
7
8      ENDDO ; monte $
9
10     ENDPROC $
11     EXEC ; silent $
12
13     ?? Writing Results to Data-Matrix
14
15
16     SAMPLE ; 1-M $
17
18     CREATE ; b1_1= 0 ; B1_2= 0 ; B1_3= 0 ; B1_4= 0 ; B1_5= 0 ; B1_6= 0 ; B1_7= 0 ; B1_8= 0
19     ; B1_9= 0 ; B1_10=0 ; B1_11=0 ; B1_12=0 ; B1_13=0 ; B1_14=0 $
20
21     NAMELIST ; VA= b1_1,B1_2,B1_3,B1_4,B1_5,B1_6,B1_7,B1_8,B1_9,B1_10,B1_11,B1_12,B1_13,B1_14 $
22
23     CREATE ; VA= BCOL $
24
25     SAMPLE ; 1-NN $
26
27
28     ?? MULTIVARIATE HECKMAN-Model with SURE (Estimator #2)
29
30     ?? MONTE-CARLO EXPERIMENT
31
32     ?? Vectors for Intermediate Output
33
34     MATRIX ; BCOL= Init(M,14,0) $
35
36     ? Renew Name-Lists
37
38     NAMELIST ; Z= ONE,Z1,Z2
39     ; X= ONE,X1,X2
40     ; Q= ONE,Q1,Q2
41     ; O= ONE,O1,O2
42     ; L1= X,M1
43     ; L2= O,M2
44     ; VA= R1,R2,R3,R4 $
45
46     PROC
47

```

```
1 DO FOR ; monte ; j= 1,M $
2
3 LABEL ; 99 $
4
5 ?? Drawing Random Error "ER"
6
7 ?? Drawing independent N(0,1) Random
8
9 CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
10 ; R3= Rnn(0,1) ; R4= Rnn(0,1) $
11
12 NAME ; VA= R1,R2,R3,R4 $
13
14 ?? Transformation to N(0,MU) distributed Random
15
16 CREA ; ER1= VA'LMU1
17 ; ER2= VA'LMU2
18 ; ER3= VA'LMU3
19 ; ER4= VA'LMU4 $
20
21 ?? Discrete Endogenous Variables
22
23 CREA ; YD1= ALF1'Z + ER1
24 ; if(YD1>0) YD1=1 ; (Else) YD1=0
25 ; YD2= ALF2'Q + ER2
26 ; if(YD2>0) YD2=1 ; (Else) YD2=0 $
27
28 ?? Continuous Endogenous Variables
29
30 CREA ; YS1=YD1*(BET1'X + ER3)
31 ; YS2=YD2*(BET2'O + ER4) $
32
33 ?? MODEL-ESTIMATION
34
35 ?? Univariate Probit-Models
36
37 PROBIT ; lhs= YD1 ; rhs= Z
38 ; start= alf1 $
39
40 MATRIX ; BP1= B $ ? Coefficient Vector
41 CREATE ; M1= N01(BP1'Z)/Phi(BP1'Z) ? Correction Terms
42 ; P1= Phi(BP1'Z) $
43
44 CALC ; exi= opt_exit+ exitcode $
45
46
47 PROBIT ; lhs= YD2 ; rhs= Q
```

C:\msys5\opt\bin\win32\bin\win32\bin
 Friday, 17 March 2006

Page 1
 14:22:00 AM

```

; start= alf2 $

MATRIX ; BP2= B $           ? Coefficient Vector
CREATE ; M2= N01(BP2'Q)/Phi(BP2'Q) ? Correction Terms
; P2= Phi(BP2'Q) $

GO TO ; 99 ; exit+ opt_exit+ exitcode # 0 $ ? Convergence Control

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-ESTIMATION

NAMELIST ; L1= X,M1
; L2= O,M2 $

CREATE ; YD12= YD1*YD2 $

?? OLS-Coefficients

MATRIX ; B1= <L1'[YD1]L1>*L1'[YD1]YS1
; B2= <L2'[YD2]L2>*L2'[YD2]YS2 $

?? Residuals

CREATE ; E11= YD1*(YS1- B1'L1)
; E22= YD2*(YS2- B2'L2) $

NAME ; VA= E11,E22 $

?? Preparation for SURE-Estimation
?? Estimation: Error-Covariances

MATRIX ; VARS= {1/NN}*VA'VA $

?? FGLS-ESTIMATION

?? Inversion and Cholesky-Factorisation of individual Covariance-Matrices

CREATE ; det= VARS(1,1)*VARS(2,2)-VARS(1,2)^2
; W11= YD1*(VARS(2,2)/(det))^0.5
; W12= YD1*YD2*(-VARS(1,2)/VARS(2,2))*(VARS(2,2)/(det))^0.5
; W22= YD2*(1/VARS(2,2))^0.5 $

CREATE ; if(YD1=1 & YD2=0) W11= YD1*(1/VARS(1,1))^0.5 $

? Data Transformation for FGLS-Estimation

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 1
04/12/2010 10:00

```
CREATE ; X111= W11*X1 ; X211= W11*X2 ; M111= W11*M1 ; YS1W= W11*YS1+ W12*YS2
; X112= W12*O1 ; X212= W12*O2 ; M212= W12*M2
; X122= W22*O1 ; X222= W22*O2 ; M222= W22*M2 ; YS2W= W22*YS2 $
```

```
? FGLS-ESTIMATION
```

```
NAME ; L1= W11,X111,X211,M111
; L12= W12,X112,X212,M212
; L2= W22,X122,X222,M222 $
```

```
MATR ; OL= L1'L1
; OR= L1'L12
; UL= L12'L1
; UR= L12'L12+ L2'L2 $
```

```
MATR ; YO= L1'YS1W
; YU= L12'YS1W+ L2'YS2W
; XW= [OL,OR/UL,UR]
; YW= [YO/YU] $
```

```
MATR ; BR= <XW'XW>*XW'YW $
```

```
? Accumulation of Results
```

```
MATR ; COEF= [BP1/BP2/BR]
; BCOL(j,*)= COEF $
```

```
ENDDO ; monte $
```

```
ENDPROC $
EXEC ; silent $
```

```
?? Writing Results to Data-Matrix
```

```
SAMPLE ; 1-M $
```

```
CREATE ; b2_1= 0 ; b2_2= 0 ; b2_3= 0 ; b2_4= 0 ; b2_5= 0 ; b2_6= 0 ; b2_7= 0 ; b2_8= 0
; b2_9= 0 ; b2_10=0 ; b2_11=0 ; b2_12=0 ; b2_13=0 ; b2_14=0 $
```

```
NAMELIST ; VA= b2_1,b2_2,b2_3,b2_4,b2_5,b2_6,b2_7,b2_8,b2_9,b2_10,b2_11,b2_12,b2_13,b2_14 $
```

```
CREATE ; VA= BCOL $
```

```
SAMPLE ; 1-NN $
```

```
?? MULTIVARITE HECKMAN with FGLS (Estimator #6)
```


1 CaseStudy_CompEconsworld_000006_1.in
 2 Friday, 17 March 2006

Page 3
 16:12:00 AM

3 ?? MONTE-CARLO EXPERIMENT

4 ?? Vectors for Intermediate Output

5 MATRIX ; BCOL= Init(M,14,0) \$

6 ? Generating Name-Lists

7
 8
 9
 10 NAMELIST ; Z= ONE,Z1,Z2
 11 ; X= ONE,X1,X2
 12 ; Q= ONE,Q1,Q2
 13 ; O= ONE,O1,O2
 14 ; L1= X,M1
 15 ; L2= O,M2
 16 ; VA= R1,R2,R3,R4 \$

17 ? Command Short-Cut

18 STRING ; ST1= r/((2*Pi)^0.5)*N01((Fit12sq)^0.5/r)/P12 \$

19
 20 PROC

21 DO FOR ; monte ; j= 1,M \$

22 LABEL ; 99 \$

23 ?? Drawing Random Error "ER"

24 ?? Drawing independent N(0,1) Random Numbers

25 CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
 26 ; R3= Rnn(0,1) ; R4= Rnn(0,1) \$

27 NAME ; VA= R1,R2,R3,R4 \$

28 ?? Transformation to N(0,MU) distributed Random Vector

29 CREA ; ER1= VA'LMU1
 30 ; ER2= VA'LMU2
 31 ; ER3= VA'LMU3
 32 ; ER4= VA'LMU4

33 ?? Discrete Endogenous Variables

34 ; YD1= ALF1'Z + ER1
 35 ; if(YD1>0) YD1=1 ; (Else) YD1=0
 36 ; YD2= ALF2'Q + ER2
 37 ; if(YD2>0) YD2=1 ; (Else) YD2=0
 38 ; YD12= YD1*YD2

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 11
11/11/2010

?? Continuous Endogenous Variables

; YS1=YD1*(BET1'X + ER3)
; YS2=YD2*(BET2'O + ER4) \$

?? ESTIMATION: BIVARIATE PROBIT MODEL

?? Bivariate Probit Model

BIVA ; lhs= YD1,YD2
; rh1= Z
; rh2= Q
; start= alf1,alf2,cor
; maxit= 25 \$

GO TO ; 99 ; opt_exit+ exitcode # 0 \$? Convergence Control

MATR ; BMP= B \$? Coefficient Vector

CALC ; r=(1-rho^2)^0.5 \$

? Mill's Ratios

CREA ; Fit1= B(1)+B(2)*Z1+B(3)*Z2
; Fit2= B(4)+B(5)*Q1+B(6)*Q2
; M1= N01(FIT1)/Phi(FIT1)
; M2= N01(FIT2)/Phi(FIT2) \$

?? PROBABILITIES

NAME ; VA= Fit1,Fit2 \$

CREA ; P1= Phi(Fit1)
; P2= Phi(Fit2)
; P12= Bvn(VA,rho)

?? Generalized Mill's Ratios

; Fit21= Fit2-rho*Fit1
; Fit12= Fit1-rho*Fit2
; Fit12sq= Fit1^2+Fit2^2-2*rho*Fit1*Fit2
; M12= (N01(Fit1)*Phi((Fit21)/r))/P12
; M21= (N01(Fit2)*Phi((Fit12)/r))/P12 \$

?? COVARIACE MATRIX REGRESSORS

Editorial Office, Dept of Economics, Warwick University, Coventry CV4 7AL, UK

```

1  CREA    ; EX1_12=  M12 +rho*M21
2          ; EX2_12=  M21 +rho*M12
3
4          ; EX11_12= 1 - (      Fit1*N01(Fit1)*Phi((Fit21)/r)
5                    + rho^2*Fit2*N01(Fit2)*Phi((Fit12)/r)
6                    - rho*"ST1"
7
8          ; EX22_12= 1 - (rho^2*Fit1*N01(Fit1)*Phi((Fit21)/r)
9                    + Fit2*N01(Fit2)*Phi((Fit12)/r)
10                   - rho*"ST1"
11
12         ; EX12_12= rho - (rho*Fit1*N01(Fit1)*Phi((Fit21)/r)
13                   + rho*Fit2*N01(Fit2)*Phi((Fit12)/r)
14                   - "ST1" $
15

```

? Centred Truncated Moments

```

16 CREA    ; VAR11_12= EX11_12- EX1_12^2
17          ; VAR22_12= EX22_12- EX2_12^2
18          ; COV12_12= EX12_12- EX1_12*EX2_12
19          ; VAR1_1= 1- Fit1*M1- M1^2
20          ; VAR2_2= 1- Fit2*M2- M2^2 $
21
22

```

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-Estimation

```

23 NAME    ; L1=  X,M1
24          ; L2=  O,M2
25          ; L21= O,M12,M21
26          ; L12= X,M12,M21 $
27

```

?? OLS-Coefficients

```

28 MATR    ; B1= <L1'[YD1]L1>*L1'[YD1]YS1
29          ; B2= <L2'[YD2]L2>*L2'[YD2]YS2
30
31          ; B21= <L21'[YD12]L21>*L21'[YD12]YS2
32          ; B12= <L12'[YD12]L12>*L12'[YD12]YS1
33
34          ; BB1=  B1(1:3) ; BB2=  B2(1:3)
35          ; BB12= B12(1:3) ; BB21= B21(1:3) $
36

```

?? Squares and Cross-Products

```

37 CREA    ; E11= YD1*(YS1- B1'L1)^2
38          ; E22= YD2*(YS2- B2'L2)^2
39          ; E12= YD12*(YS1- B12'L12)*(YS2- B21'L21) $
40

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 11
11/12/2010 10:11:00 AM

?? Preparation: GLS-Estimation

NAME ; VA= ONE,VAR1_1 \$
MATR ; BB= <VA'[YD1]VA>*VA'[YD1]E11 \$
CREA ; V11= BB'VA \$

NAME ; VA= ONE,VAR2_2 \$
MATR ; BB= <VA'[YD2]VA>*VA'[YD2]E22 \$
CREA ; V22= BB'VA \$

NAME ; VA= ONE,VAR11_12,VAR22_12,COV12_12 \$
MATR ; BB= <VA'[YD12]VA>*VA'[YD12]E12 \$
CREA ; V12= BB'VA
+((B1'L1- BB1'X)- (B12'L12- BB12'X))
*((B2'L2- BB2'O)- (B21'L21- BB21'O)) \$

?? FGLS-ESTIMATION

?? Inversion and Cholesky-Factorisation of individual Covariance-Matrices

CREA ; det= V11*V22-V12^2
; if(YD1=1 & YD2=1 & det>0) |
W11= (V22/det)^0.5
; W12= (-V12/V22)*(V22/det)^0.5
; W22= (1/V22)^0.5
; (Else) | W11= 0 ; W22= 0 ; W12=0 \$
CREA ; if(YD1=1 & YD2=0) W11= (1/V11)^0.5
; if(YD1=0 & YD2=1) W22= (1/V22)^0.5 \$

? Checking for Non-Definiteness

? CREA ; di= 0 ; if(YD1=1 & YD2=1 & det<=0) di= 1 \$
? CALC ; din= din+ xbr(di) \$

? Data Transformation for FGLS-Estimation

CREA ; X111= W11*X1 ; X211= W11*X2 ; M111= W11*M1
; YS1W= W11*YS1+ W12*YS2
; X112= W12*O1 ; X212= W12*O2 ; M212= W12*M2
; X122= W22*O1 ; X222= W22*O2 ; M222= W22*M2
; YS2W= W22*YS2 \$

? FGLS-Estimation

NAME ; L1= W11,X111,X211,M111
; L12= W12,X112,X212,M212

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Passing Copy@Economics@Warwick_001006.Lim
Friday, 17 March 2006

Page 44
04/12/2006 10:00 AM

?? Vectors for Intermediate Output

MATRIX ; BCOL= Init(M,16,0) \$

? Generating Name-Lists

NAMELIST ; Z= ONE,Z1,Z2
; X= ONE,X1,X2
; Q= ONE,Q1,Q2
; O= ONE,O1,O2
; VA= R1,R2,R3,R4 \$

? Command Short-Cut

STRING ; ST1= r/((2*Pi)^0.5)*N01((Fit12sq)^0.5/r)/P12 \$

PROC

DO FOR ; monte ; j= 1,M \$

LABEL ; 99 \$

?? Drawing Random Error "ER"

?? Drawing independent N(0,1) Random Numbers

CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
; R3= Rnn(0,1) ; R4= Rnn(0,1) \$

NAME ; VA= R1,R2,R3,R4 \$

?? Transformation to N(0,MU) distributed Random Vector

CREA ; ER1= VA'LMU1
; ER2= VA'LMU2
; ER3= VA'LMU3
; ER4= VA'LMU4

?? Discrete Endogenous Variables

; YD1= ALF1'Z + ER1
; if(YD1>0) YD1=1 ; (Else) YD1=0
; YD2= ALF2'Q + ER2
; if(YD2>0) YD2=1 ; (Else) YD2=0
; YD12= YD1*YD2

?? Continuous Endogenous Variables

Editorial Office, Dept of Economics, Warwick University, Coventry CV4 7AL, UK

```

; YS1=YD1*(BET1'X + ER3)
; YS2=YD2*(BET2'O + ER4) $

?? ESTIMATION: BIVARIATE PROBIT MODEL

?? Bivariate Probit Model

BIVA ; lhs= YD1,YD2
; rh1= Z
; rh2= Q
; start= alf1,alf2,cor
; maxit= 25 $

GO TO ; 99 ; opt_exit+ exitcode # 0 $ ? Convergence Control

MATR ; BMP= B $ ? Coefficient Vector

CREA ; mrho= rho*(2*YD1-1)*(2*YD2-1) ; r=(1-mrho^2)^0.5 $

? Predictions

CREA ; Fit1= (B(1)+B(2)*Z1+B(3)*Z2)*(2*YD1-1)
; Fit2= (B(4)+B(5)*Q1+B(6)*Q2)*(2*YD2-1) $

?? PROBABILITIES

NAME ; VA= Fit1,Fit2 $

CREA ; P1= Phi(Fit1)
; P2= Phi(Fit2)
; P12= Bvn(VA,mrho)

?? Generalized Mill's Ratios

; Fit21= Fit2-mrho*Fit1
; Fit12= Fit1-mrho*Fit2

; M12= (2*YD1-1)*(N01(Fit1)*Phi((Fit21)/r))/P12
; M21= (2*YD2-1)*(N01(Fit2)*Phi((Fit12)/r))/P12 $

?? Weighted Variables

CREATE ; X1Y1= X1*YD1 ; X2Y1= X2*YD1 ; M12Y1= M12*YD1 ; M21Y1= M21*YD1
; X1Y2= O1*YD2 ; X2Y2= O2*YD2 ; M21Y2= M21*YD2 ; M12Y2= M12*YD2 $

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 46
14/12/2010 10:00

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-Estimation

NAME ; L1= YD1,X1Y1,X2Y1,M12Y1,M21Y1
; L2= YD2,X1Y2,X2Y2,M21Y2,M12Y2 \$

MATR ; OL= L1'L1
; OR= Init(5,5,0)
; UL= Init(5,5,0)
; UR= L2'L2

; YO= L1'YS1
; YU= L2'YS2
; XW= [OL,OR/UL,UR]
; YW= [YO/YU]

; BR= <XW'XW>*XW'YW \$

? Accumulation of Results

MATR ; COEF= [BMP/BR]
; BCOL(j,*)= COEF \$

ENDDO ; monte \$

ENDPROC \$
EXEC ; silent \$

?? Writing Results to Data-Matrix

SAMPLE ; 1-M \$

CREATE ; b10_1= 0 ; b10_2= 0 ; b10_3= 0 ; b10_4= 0 ; b10_5= 0 ; b10_6= 0 ; b10_7= 0 ; b10_8= 0
; b10_9= 0 ; b10_10=0 ; b10_11=0 ; b10_12=0 ; b10_13=0 ; b10_14=0 ; b10_15=0 ; b10_16=0 \$

NAMelist ; VA= b10_1,b10_2,b10_3,b10_4,b10_5,b10_6,b10_7,b10_8,b10_9,b10_10,b10_11,b10_12,b10_13,b10_14,b10_15,b10_16 \$

CREATE ; VA= BCOL \$

SAMPLE ; 1-NN \$

??

?? MULTIVARITE HECKMAN with Full Correction and SURE (Estimator # 11)

?? MONTE-CARLO EXPERIMENT

?? Vectors for Intermediate Output

1 ansys_001@econswarwick.ac.uk
2 Friday, 17 March 2006

Page 11
14:22:00 AM

3 MATRIX ; BCOL= Init(M,16,0) \$

4
5
6 ? Generating Name-Lists

7 NAMELIST ; Z= ONE,Z1,Z2
8 ; X= ONE,X1,X2
9 ; Q= ONE,Q1,Q2
10 ; O= ONE,O1,O2
11 ; VA= R1,R2,R3,R4 \$

12
13 ? Command Short-Cut

14 STRING ; ST1= r/((2*Pi)^0.5)*N01((Fit12sq)^0.5/r)/P12 \$

15
16
17 PROC

18 DO FOR ; monte ; j= 1,M \$

19 ?? LABEL ; 99 \$

20 ?? Drawing Random Error "ER"

21 ?? Drawing independent N(0,1) Random Numbers

22 CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
23 ; R3= Rnn(0,1) ; R4= Rnn(0,1) \$

24 NAME ; VA= R1,R2,R3,R4 \$

25 ?? Transformation to N(0,MU) distributed Random Vector

26 CREA ; ER1= VA'LMU1
27 ; ER2= VA'LMU2
28 ; ER3= VA'LMU3
29 ; ER4= VA'LMU4

30 ?? Discrete Endogenous Variables

31 ; YD1= ALF1'Z + ER1
32 ; if(YD1>0) YD1=1 ; (Else) YD1=0
33 ; YD2= ALF2'Q + ER2
34 ; if(YD2>0) YD2=1 ; (Else) YD2=0
35 ; YD12= YD1*YD2

36 ?? Continuous Endogenous Variables

37 ; YS1=YD1*(BET1'X + ER3)

38
39
40
41
42
43
44
45 **Editorial Office, Dept of Economics, Warwick University, Coventry CV4 7AL, UK**
46
47

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 11
11/11/2006

```

; YS2=YD2*(BET2'O + ER4) $

?? ESTIMATION: BIVARIATE PROBIT MODEL

?? Bivariate Probit Model

BIVA ; lhs= YD1,YD2
      ; rh1= Z
      ; rh2= Q
      ; start= alf1,alf2,cor
      ; maxit= 25 $

GO TO ; 99 ; opt_exit+ exitcode # 0 $ ? Convergence Control

MATR ; BMP= B $ ? Coefficient Vector

CREA ; mrho= rho*(2*YD1-1)*(2*YD2-1) ; r=(1-mrho^2)^0.5 $

? Predictions

CREA ; Fit1= (BMP(1)+BMP(2)*Z1+BMP(3)*Z2)*(2*YD1-1)
      ; Fit2= (BMP(4)+BMP(5)*Q1+BMP(6)*Q2)*(2*YD2-1) $

?? PROBABILITIES

NAME ; VA= Fit1,Fit2 $

CREA ; P1= Phi(Fit1)
      ; P2= Phi(Fit2)
      ; P12= Bvn(VA,mrho)

?? Generalized Mill's Ratios

      ; Fit21= Fit2-mrho*Fit1
      ; Fit12= Fit1-mrho*Fit2

      ; M12= (2*YD1-1)*(N01(Fit1)*Phi((Fit21)/r))/P12
      ; M21= (2*YD2-1)*(N01(Fit2)*Phi((Fit12)/r))/P12 $

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-ESTIMATION

NAMELIST ; L1= X,M12,M21
          ; L2= O,M21,M12 $

CREATE ; YD12= YD1*YD2 $

```

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

?? OLS-Coefficients

```
MATRIX ; B1= <L1'[YD1]L1>*L1'[YD1]YS1
        ; B2= <L2'[YD2]L2>*L2'[YD2]YS2 $
```

?? Residuals

```
CREATE ; E11= YD1*(YS1- B1'L1)
        ; E22= YD2*(YS2- B2'L2) $
```

```
NAME ; VA= E11,E22 $
```

?? Preparation for SURE-Estimation

?? Estimation: Error-Covariances

```
MATRIX ; VARS= {1/NN}*VA'VA $
```

?? FGLS-ESTIMATION

?? Inversion and Cholesky-Factorisation of individual Covariance-Matrices

```
CREATE ; det= VARS(1,1)*VARS(2,2)-VARS(1,2)^2
        ; W11= YD1*(VARS(2,2)/(det))^0.5
        ; W12= YD1*YD2*(-VARS(1,2)/VARS(2,2))*(VARS(2,2)/(det))^0.5
        ; W22= YD2*(1/VARS(2,2))^0.5 $
```

```
CREATE ; if(YD1=1 & YD2=0) W11= YD1*(1/VARS(1,1))^0.5 $
```

? Data Transformation for FGLS-Estimation

```
CREATE ; X111= W11*X1 ; X211= W11*X2 ; M1211= W11*M12 ; M2111= W11*M21 ; YS1W= W11*YS1+ W12*YS2
        ; X112= W12*O1 ; X212= W12*O2 ; M2112= W12*M21 ; M1212= W12*M12
        ; X122= W22*O1 ; X222= W22*O2 ; M2122= W22*M21 ; M1222= W22*M12 ; YS2W= W22*YS2 $
```

? FGLS-ESTIMATION

```
NAME ; L1= W11,X111,X211,M1211,M2111
        ; L12= W12,X112,X212,M2112,M1212
        ; L2= W22,X122,X222,M2122,M1222 $
```

```
MATR ; OL= L1'L1
        ; OR= L1'L12
        ; UL= L12'L1
        ; UR= L12'L12+ L2'L2
```

```
; YO= L1'YS1W
```


Working Paper in Economics and Statistics, 2010/10
 Friday, 17 March 2010

Page 11
 14:22:00

```

3           ; VA= R1,R2,R3,R4 $
4
5 ? Command Short-Cut
6
7 STRING ; ST1= r/((2*Pi)^0.5)*N01((Fit12sq)^0.5/r)/P12 $
8
9 PROC $
10     DO FOR ; monte ; j= 1,M $
11         LABEL ; 99 $
12             ?? Drawing Random Error "ER"
13             ?? Drawing independent N(0,1) Random Numbers
14
15             CREA ; R1= Rnn(0,1) ; R2= Rnn(0,1)
16                 ; R3= Rnn(0,1) ; R4= Rnn(0,1) $
17
18             NAME ; VA= R1,R2,R3,R4 $
19
20             ?? Transformation to N(0,MU) distributed Random Vector
21
22             CREA ; ER1= VA'LMU1
23                 ; ER2= VA'LMU2
24                 ; ER3= VA'LMU3
25                 ; ER4= VA'LMU4
26
27             ?? Discrete Endogenous Variables
28                 ; YD1= ALF1'Z + ER1
29                 ; if(YD1>0) YD1=1 ; (Else) YD1=0
30                 ; YD2= ALF2'Q + ER2
31                 ; if(YD2>0) YD2=1 ; (Else) YD2=0
32                 ; YD12= YD1*YD2
33
34             ?? Continuous Endogenous Variables
35                 ; YS1=YD1*(BET1'X + ER3)
36                 ; YS2=YD2*(BET2'O + ER4) $
37
38             ?? ESTIMATION: BIVARIATE PROBIT MODEL
39             ?? Bivariate Probit Model
40
41             BIVA ; lhs= YD1,YD2
42                 ; rh1= Z
43                 ; rh2= Q
44                 ; start= alf1,alf2,cor

```

Classiq_CompEconWorkBook_001006.Lim
Friday, 17 March 2006

Page 11
14:22:00 AM

```

; maxit= 25 $

GO TO ; 99 ; opt_exit+ exitcode # 0 $ ? Convergence Control

MATR ; BMP= B $ ? Coefficient Vector

CREATE ; mrho= rho*(2*YD1-1)*(2*YD2-1) ; r=(1-mrho^2)^0.5 $

? Mill's Ratios

CREA ; Fit1= (BMP(1)+BMP(2)*Z1+BMP(3)*Z2)*(2*YD1-1)
; Fit2= (BMP(4)+BMP(5)*Q1+BMP(6)*Q2)*(2*YD2-1)
; M1= N01(FIT1)/Phi(FIT1)
; M2= N01(FIT2)/Phi(FIT2) $

?? PROBABILITIES

NAME ; VA= Fit1,Fit2 $

CREA ; P1= Phi(Fit1)
; P2= Phi(Fit2)
; P12= Bvn(VA,mrho)

?? Generalized Mill's Ratios

; Fit21= Fit2-mrho*Fit1
; Fit12= Fit1-mrho*Fit2
; Fit12sq= Fit1^2+Fit2^2-2*mrho*Fit1*Fit2
; M12= (2*YD1-1)*(N01(Fit1)*Phi((Fit21)/r))/P12
; M21= (2*YD2-1)*(N01(Fit2)*Phi((Fit12)/r))/P12 $

?? COVARIACE MATRIX REGRESSORS

CREA ; EX1_12= M12 +mrho*M21
; EX2_12= M21 +mrho*M12

; EX11_12= 1 -( Fit1*N01(Fit1)*Phi((Fit21)/r)
+ mrho^2*Fit2*N01(Fit2)*Phi((Fit12)/r)
- mrho*"ST1"

; EX22_12= 1 -(mrho^2*Fit1*N01(Fit1)*Phi((Fit21)/r)
+ Fit2*N01(Fit2)*Phi((Fit12)/r)
- mrho*"ST1"

; EX12_12= cor -(mrho*Fit1*N01(Fit1)*Phi((Fit21)/r)
+ mrho*Fit2*N01(Fit2)*Phi((Fit12)/r)
- "ST1" $

```

? Centred Truncated Moments

```
CREA ; VAR11_12= EX11_12- EX1_12^2
      ; VAR22_12= EX22_12- EX2_12^2
      ; COV12_12= EX12_12- EX1_12*EX2_12 $
```

?? ESTIMATION: CONTINUOUS MODEL

?? OLS-Estimation

```
NAME ; L12= X,M12,M21
      ; L21= O,M12,M21 $
```

?? OLS-Coefficients

```
MATR ; B21= <L21'[YD12]L21>*L21'[YD12]YS2
      ; B12= <L12'[YD12]L12>*L12'[YD12]YS1
      ; BB12= B12(1:3) ; BB21= B21(1:3) $
```

?? Squares and Cross-Products

```
CREA ; E11= YD1*(YS1- B12'L12)^2
      ; E22= YD2*(YS2- B21'L21)^2
      ; E12= YD12*(YS1- B12'L12)*(YS2- B21'L21) $
```

?? Preparation: GLS-Estimation

```
NAME ; VA= ONE,VAR11_12,VAR22_12,COV12_12 $
MATR ; BB= <VA'[YD1]VA>*VA'[YD1]E11 $
CREA ; V11= BB'VA $
```

```
MATR ; BB= <VA'[YD2]VA>*VA'[YD2]E22 $
CREA ; V22= BB'VA $
```

```
MATR ; BB= <VA'[YD12]VA>*VA'[YD12]E12 $
CREA ; V12= BB'VA
      ; det= V11*V22-V12^2
```

?? Checking for Non-Definiteness

```
; V11D= YD1*V11 ; V22D= YD2*V22 ; detd= YD12*det $
```

```
CALC ; ch1= min(V11D) ; ch2= min(V22D) ; ch3= min(detd) $
```

```
GO TO ; 99 ; ch1<0 | ch2<0 | ch3<0 $ ? Control for Non-Definiteness
```

?? FGLS-ESTIMATION

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Page 11
14:12:00 AM

?? Inversion and Cholesky-Factorisation of individual Covariance-Matrices

```
CREA ; if(YD12=1) |
      W11= (V22/det)^0.5
      ; W12= (-V12/V22)*(V22/det)^0.5
      ; W22= (1/V22)^0.5
      ; (Else) | W11= 0 ; W22= 0 ; W12=0 $

CREA ; if(YD1=1 & YD2=0) W11= (1/V11)^0.5
      ; if(YD1=0 & YD2=1) W22= (1/V22)^0.5 $
```

? Data Transformation for FGLS-Estimation

```
CREATE ; X111= W11*X1 ; X211= W11*X2 ; M1211= W11*M12 ; M2111= W11*M21 ; YS1W= W11*YS1+ W12*YS2
      ; X112= W12*O1 ; X212= W12*O2 ; M2112= W12*M21 ; M1212= W12*M12
      ; X122= W22*O1 ; X222= W22*O2 ; M2122= W22*M21 ; M1222= W22*M12 ; YS2W= W22*YS2 $
```

? FGLS-ESTIMATION

```
NAME ; L1= W11,X111,X211,M1211,M2111
      ; L12= W12,X112,X212,M2112,M1212
      ; L2= W22,X122,X222,M2122,M1222 $
```

```
MATR ; OL= L1'L1
      ; OR= L1'L12
      ; UL= L12'L1
      ; UR= L12'L12+ L2'L2

      ; YO= L1'YS1W
      ; YU= L12'YS1W+ L2'YS2W
      ; XW= [OL,OR/UL,UR]
      ; YW= [YO/YU]

      ; BR= <XW'XW>*XW'YW $
```

? Accumulation of Results

```
MATR ; COEF= [BMP/BR]
      ; BCOL(j,*)= COEF $
```

ENDDO ; monte \$

ENDPROC \$
EXEC ; silent \$

?? Writing Results to Data-Matrix

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

Casey_Comp@Warwick.ac.uk_Miles_001061.in
Friday, 17 March 2006

Page 15
04/12/2010 10:00

SAMPLE ; 1-M \$

CREATE ; b12_1= 0 ; b12_2= 0 ; b12_3= 0 ; b12_4= 0 ; b12_5= 0 ; b12_6= 0 ; b12_7= 0 ; b12_8= 0
; b12_9= 0 ; b12_10=0 ; b12_11=0 ; b12_12=0 ; b12_13=0 ; b12_14=0 ; b12_15=0 ; b12_16=0 \$

NAMELIST ; VA= b12_1,b12_2,b12_3,b12_4,b12_5,b12_6,b12_7,b12_8,b12_9,b12_10,b12_11,b12_12,b12_13,b12_14,b12_15,b12_16 \$

CREATE ; VA= BCOL \$

SAMPLE ; 1-NN \$

For Peer Review

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Peer Review