

Disentangling the effects of trait and state worry on error-related brain activity: Results from a randomized controlled trial using worry manipulations

Härpfer, Kai; Carsten, Hannes Per; Löwisch, Kim; Westermann, Nele; Riesel, Anja

Veröffentlichungsversion / Published Version
Zeitschriftenartikel / journal article

Empfohlene Zitierung / Suggested Citation:

Härpfer, K., Carsten, H. P., Löwisch, K., Westermann, N., & Riesel, A. (2022). Disentangling the effects of trait and state worry on error-related brain activity: Results from a randomized controlled trial using worry manipulations. *Psychophysiology*, 59(9), 1-20. <https://doi.org/10.1111/psyp.14055>

Nutzungsbedingungen:

Dieser Text wird unter einer CC BY Lizenz (Namensnennung) zur Verfügung gestellt. Nähere Auskünfte zu den CC-Lizenzen finden Sie hier:
<https://creativecommons.org/licenses/by/4.0/deed.de>

Terms of use:

This document is made available under a CC BY Licence (Attribution). For more Information see:
<https://creativecommons.org/licenses/by/4.0>

Disentangling the effects of trait and state worry on error-related brain activity: Results from a randomized controlled trial using worry manipulations

Kai Härpfer  | Hannes Per Carsten  | Kim Löwisch  | Nele Westermann  | Anja Riesel 

Department of Psychology, University of Hamburg, Hamburg, Germany

Correspondence

Kai Härpfer, Department of Psychology, University of Hamburg, Von-Melle-Park 11, 20146 Hamburg, Germany.
Email: kai.haerpfer@uni-hamburg.de

Funding information

This research was supported by the German Research Foundation (grant awarded to Anja Riesel DFG-Grant RI-2853/2-1). Parts of the data have been presented previously on the SPR Annual Meeting 2021.

Abstract

Enhanced amplitudes of the error-related negativity (ERN) have been suggested to be a transdiagnostic neural risk marker for internalizing psychopathology. Previous studies propose worry to be an underlying mechanism driving the association between enhanced ERN and anxiety. The present preregistered study focused on disentangling possible effects of trait and state worry on the ERN by utilizing a cross sectional observational and a longitudinal randomized controlled experimental design. To this end, we examined the ERN of $n = 90$ students during a flanker task (T0), which were then randomly assigned to one of three groups (worry induction, worry reduction, passive control group). Following the intervention, participants performed another flanker task (T1) to determine potential alterations of their ERN. Manipulation checks revealed that compared to the control group, state worry increased in the induction but also in the reduction group. ERN amplitudes did not vary as a function of state worry. An association of trait worry with larger ERN amplitudes was only observed in females. Furthermore, we found larger ERN amplitudes in participants with a current or lifetime diagnosis of internalizing disorders. In summary, our findings suggest that the ERN seems to be insensitive to variations in state worry, but that an elevated ERN is associated with the trait-like tendency to worry and internalizing psychopathology, which is consistent with the notion that the ERN likely represents a trait-like neural risk associated with anxiety.

KEYWORDS

anxiety, anxious apprehension, EEG, ERN, error monitoring, worry

1 | INTRODUCTION

Goal-directed behavior is a fundamental ability allowing humans to adjust to their environment. Cognitive control, especially performance monitoring, plays a crucial role for goal-directed behavior. By monitoring errors (i.e., potential harm), humans can improve upcoming behavior through cognitive, motivational, and behavioral adjustments (Botvinick et al., 2001; Cavanagh & Shackman, 2015; Proudfit et al., 2013; Simons, 2010; Weinberg et al., 2012).

A commonly studied event-related potential (ERP) associated with performance monitoring is the error-related negativity (ERN; Gehring et al., 1993); or originally called the error negativity (N_E ; Falkenstein et al., 1991). The ERN is assumed to constitute the neural representative of error monitoring and is observable as a fronto-central negative peak approx. 50 ms after committing an error. The monitoring of correct responses elicits a similarly timed but smaller negative peak after a correct response: the correct-response negativity (CRN; Vidal et al., 2003, 2000). Various hypotheses have been postulated about the functional role of the ERN, ranging from signaling the mismatch between representations of required and actual responses (Falkenstein et al., 1991; Gehring et al., 1993), over indicating the conflict of two simultaneously active response tendencies (Botvinick et al., 2001; Yeung et al., 2004), to allowing reinforcement learning to modify performance on a task at hand by signaling whenever an outcome was not as predicted (Holroyd & Coles, 2002). All of these hypotheses suggest that the ERN is crucial for cognitive control and a prerequisite to adjusting behavior to the requirements of the task and improving future performance.

Regarding the clinical utility of error monitoring (Hajcak et al., 2019), increased or decreased ERN amplitudes have been discussed in the literature as an endophenotype for the development and maintenance of psychopathological symptoms (e.g., Manoach & Agam, 2013; Olvet & Hajcak, 2008; Riesel, Klawohn, et al., 2019). An endophenotype is defined as a measurable (mostly, but not necessarily, biological) component on the complex pathway between the genotype and a psychopathological phenotype, informative of the specific mechanisms that lead to a complex mental disorder (Gottesman & Gould, 2003). A marker qualifies as an endophenotype (a) when it is associated with the illness, (b) when it is heritable, (c) when it is primarily state-independent, (d) when the endophenotype and the illness co-segregate within families, and (e) when the endophenotype is also more prevalent in nonaffected family members compared to individuals of the general population (Gottesman & Gould, 2003).

In fact, there is convincing meta-analytical evidence for an association of the ERN with psychopathology (criterion

a): ERN variations were found along the lines of internalizing and externalizing mental disorders (Lutz et al., 2021; Pasion & Barbosa, 2019) with enhanced ERN amplitudes for anxiety (Moser et al., 2013; Saunders & Inzlicht, 2020) and obsessive-compulsive disorders (Riesel, 2019) on the one hand, and attenuated ERN amplitudes for substance use disorder (Luijten et al., 2014) and attention deficit hyperactivity disorder (Shiels & Hawk, 2010) on the other hand. The ERN also fulfills criterion (b), as it has been found to be heritable from one generation to the other (Anokhin et al., 2008; Suor et al., 2021), and criterion (c), since a successful cognitive-behavioral therapy decreasing psychopathological symptoms has no effect on the ERN (Gorka et al., 2018; Hajcak et al., 2008; Kujawa et al., 2016; Ladouceur et al., 2018; Riesel et al., 2015). Lastly, corresponding alterations of the ERN have also been found in unaffected individuals with a family history of anxiety disorders, obsessive-compulsive disorders, and substance use disorder (Carrasco et al., 2013; Riesel et al., 2011; Riesel, Klawohn, et al., 2019), implying co-segregation within families (criterion d) and higher rates in first-degree relatives (criterion e). In addition, enhanced ERN amplitudes predict the onset of anxiety disorders (Meyer et al., 2015), all together supporting the notion that increased ERN amplitudes represent a trait-like neural risk marker or endophenotype for anxiety.

As mentioned above, the endophenotype approach aims at identifying specific mechanisms that lead to psychopathology. Unfortunately, the identification of mechanisms leading to pathological anxiety is complicated by the complex and heterogeneous nature of anxiety that likely overshadows specific associations between neural functioning and psychopathology. Regarding different dimensions of anxiety, the link between anxiety and elevated ERN amplitudes seems to be driven by worry as opposed to anxious arousal. This is supported by previous studies (Hajcak et al., 2003; Lin et al., 2015; Moran et al., 2012; Moser et al., 2012; but see Härpfer, Carsten, Spychalski, et al., 2020) as well as meta-analyses (Moser et al., 2013; Saunders & Inzlicht, 2020). Specifically, enhanced error monitoring in clinical populations such as generalized anxiety disorder (GAD; Meyer et al., 2012; Weinberg et al., 2010), social anxiety disorder (SAD; Endrass et al., 2014), health anxiety (Riesel et al., 2017), or obsessive-compulsive disorder (OCD; see Mathews et al., 2012; Riesel, 2019, for reviews and a meta-analysis), has been assumed to be related to transdiagnostically shared worry symptoms. Studies of healthy subjects and meta-analyses suggest that this relationship is evident across the full spectrum of worry symptoms including subclinical individuals. However, this assumption has been challenged in more recent studies, which tend to suggest that the association may be stronger in clinical

populations (Saunders & Inzlicht, 2020). In addition, gender seems to be an important moderator, and especially women show the expected association between worry and ERN (Moser et al., 2016).

Although the ERN is often discussed as an endophenotype, studies that experimentally manipulated state affect (e.g., Nigbur & Ullsperger, 2020; Wiswede, Münte, Goschke, et al., 2009), attentional biases (Klawohn, Hajcak, et al., 2020; Nelson et al., 2015), or the consequences of an error (i.e., punishment; Meyer & Gawłowska, 2017; Riesel et al., 2012) found that the ERN is susceptible for intra-individual variation. Utilizing the strengths of a causal intervention study, a recent approach that focused on the experimental reduction of worry (Schroder et al., 2018) showed that emotional expressive writing was associated with attenuated ERN amplitudes, which points to state worry as an affective state associated with ERN variations. This central and important study used a between group design, limiting conclusions because of possible confounds introduced by preexisting between-group variations. Thus, longitudinal within-between comparisons are important and promising to extend this line of research.

In summary, a variety of findings point to an association between ERN and worry. Based on this, the compensatory error-monitoring hypothesis (CEMH; Moser et al., 2013) postulates that anxious individuals need to employ compensatory effort, as reflected by an increased ERN, in order to overcome processing inefficiency that is caused by the distracting effects of worry on working memory capacity. As a result, the compensating effort leads to comparable levels of task performance. Another influential approach (Proudfit et al., 2013) assumes that enhanced error monitoring of anxious individuals is caused by a pronounced trait-like sensitivity to uncertain threats (e.g., errors): This threat sensitivity temporally increases defensive motivation in an uncertain and potentially threatening situation, making errors motivationally more relevant and leading to greater error monitoring. In this view, worrying is a by-product that has developed as a maladaptive coping strategy of anxious individuals associated with heightened threat sensitivity. In accordance with the endophenotype approach, the authors argue that ERN variations are due to differences in threat sensitivity—a stable trait—not due to state-dependent temporarily efforts to compensate for the distracting effects of worries. Overall, both approaches converge on the idea that anxiety is associated with an increased ERN, although they differ in their assumptions on the underlying mechanisms of this relationship.

This preregistered study (Härpfer et al., 2020) aimed at disentangling the relationships between trait and state worry and error-related brain activity by utilizing both a cross sectional observational design (similar to many previous studies) as well as longitudinal randomized controlled

experimental design manipulating state worry (allowing causal inferences). We wanted to examine whether worry interventions cause alterations in neural signals of performance monitoring. To this end, we assessed the baseline ERN of 90 participants (T0), which were then randomly assigned to one of three groups (two experimental groups with either a worry induction or reduction; one passive control group with no worry intervention). Following the intervention, participants performed another flanker task to determine potential alterations of their ERN (T1). Our overall research question targeted the relationship between the ERN and trait worry as well as state worry, i.e., whether this link can be found in both cross sectional and longitudinal comparisons. As preregistered, we expected that trait worry would be associated with increased ERN amplitudes at T0 and that state worry would cause alterations in ERN amplitudes in such way that, relative to the control group, ERN amplitudes between T0 and T1 would increase in the worry induction group and decrease in the worry reduction group. Likewise, we exploratorily investigated associations of trait and state worry and the CRN cross-sectionally and longitudinally. In a first meta-analysis, there was no evidence for a link between anxiety and the CRN (Moser et al., 2013). In contrast, a more recent meta-analysis found a small but significant association of anxiety and the CRN, but this was not moderated by the type of anxiety, such as worry (Saunders & Inzlicht, 2020).

2 | METHOD

2.1 | Participants

When preregistering the hypotheses and methods of our study (Härpfer et al., 2020), an a priori sample size calculation was conducted using G*Power, version 3.1.9.7 (Faul et al., 2009, 2007). Based on the results of a previous study (Schroder et al., 2018), we assume to find medium-sized effects of the worry interventions. Paired comparisons of subgroups (two-sided dependent *t*-tests) can detect medium-sized effects (Cohen's $d > 0.60$) with a sample size of $n = 24$ per group, a power of 80%, and an alpha of 0.05 (Cohen, 1992). Specifications of the sample size calculation can be found in the supplementary materials (Figure S1). As preregistered, participants were recruited until $n = 30$ complete and evaluable data sets per group were collected ($n = 4$ were excluded and replaced, for details see section 'Electrophysiological Recording and Processing'). In light of the mixed previous findings, this enabled detecting possible smaller effects. Therefore, our final sample consisted of $N = 90$ right-handed university students (66 identified as

female) aged 18 to 30 years ($M = 23.50$, $SD = 3.12$). They received either course credit or monetary compensation for their participation.

Participants were required to speak German as a native language, to have normal or corrected-to-normal vision, and to be able to provide written informed consent. Exclusion criteria for all subjects included a history of any neurological disorder, current or lifetime diagnosis of a substance-related disorder, schizophrenia spectrum disorder, bipolar disorder, and use of benzodiazepines during the last week or of neuroleptic medication during the last three months. At the time of participation, $n = 15$ participants were currently medicated with at least one drug including oral contraceptives ($n = 7$), antidepressants ($n = 4$), dermatological drugs against acne ($n = 3$), and thyroid hormones ($n = 2$).

Regarding clinical status, $n = 14$ participants had a current or lifetime diagnosis for at least one mental disorder including a major depressive episode ($n = 5$ lifetime), anorexia nervosa ($n = 3$ lifetime), bulimia nervosa ($n = 3$ lifetime), specific phobia ($n = 2$ current, $n = 1$ lifetime), panic disorder ($n = 2$ lifetime), obsessive-compulsive disorder ($n = 1$ lifetime), pain disorder ($n = 1$ lifetime), and posttraumatic stress disorder ($n = 1$ lifetime). Note, that information on clinical status was missing for seven participants.

2.2 | Procedure

Participants received verbal and written information of the objectives and methods of the study and gave written informed consent. Mental disorders were assessed by trained personnel using the Structured Clinical Interview for DSM-5—clinical version (SCID-5-CV; Beesdo-Baum et al., 2019; First et al., 2016). During the laboratory assessment (Figure 1), participants were asked to identify at least three worry topics with high personal relevance that could be used at a later point in time. They were given a list of possible content domains as examples (Arch & Craske, 2006; Boehnke et al., 1998) including social relations, achievement/work, money/economics, health, and safety, but they could also write down any other ideographic worry topic. This identification procedure was adapted from previous worry intervention studies (Arch & Craske, 2006; Oathes et al., 2008; Vasey & Borkovec, 1992; Verkuil et al., 2009).

After that, participants completed several questionnaires, performed a flanker task (T0), and had a four-minute resting state assessment (T0). The resting state served for another research question and results will be reported elsewhere. Next, participants were randomly assigned to one of three groups (induction, reduction,

control) and parallelized across groups regarding gender. This randomization procedure ensured a balanced design with equally sized groups. The experimental groups received either an eight-minute worry induction or reduction; the passive control group did not receive any intervention and paused for an equivalent amount of time. The interventions were informed by previous literature. We aimed at selecting interventions that were as potent and standardized as possible. Regarding the length of the interventions, we considered eight minutes as the best tradeoff between sufficiently inducing worries in the induction group, yet preventing participants to start habituating, and reducing worry effectively in the reduction group by allowing them enough time to reflect.

The worry induction was consistent with previous studies using a classical induction paradigm to create a worrisome and ruminative state (e.g., Arch & Craske, 2006; Borkovec & Inz, 1990; Fisher & Newman, 2013; Lyonfields et al., 1995; McLaughlin et al., 2007; Oathes et al., 2008; Ray et al., 2009; Ruscio & Borkovec, 2004; Thayer et al., 1996; Vasey & Borkovec, 1992; Verkuil et al., 2009). In accordance with the Catastrophizing Interview Technique (Vasey & Borkovec, 1992), participants were instructed to worry as intensively as they can until the experimenter asked them to stop. Participants should think about worst-case scenarios, the consequences for themselves as well as for significant others, and how badly they would feel if their worries became reality. They were allowed to switch back and forth between worry topics to facilitate rumination; but they were instructed to return to the chosen worry topics if their thoughts drifted away.

The reduction paradigm was also based on previous research on emotional expressive writing (e.g., Baddeley & Pennebaker, 2011; Gortner et al., 2006; Pennebaker & Beall, 1986; Pennebaker & Francis, 1996; Ramirez & Beilock, 2011; Sayer et al., 2015; Schroder et al., 2018). Participants were asked to write as openly as possible about their thoughts and feelings regarding their worry topics until the experimenter asked them to stop. Participants were encouraged to explore their thoughts and feelings in a completely non-judgmental manner and they were informed that they could keep their essay to ensure confidentiality. Subsequently, all participants performed another flanker task (T1), followed by a two minute booster session of the intervention or pause, and another four-minute resting state (T1).

Manipulation checks were implemented at several points throughout the assessment (i.e., before the T0 and T1 flanker, before the T0 and T1 resting state, before the intervention/pause, and after the T1 resting state) in order to track intraindividual changes of participants'




FIGURE 1 Flowchart of the study procedure. Boxes with surrounding dashed lines were part of another research question whose results will be presented elsewhere. Duration of each task is approximative. State measurements throughout the study ensured the tracking of fluctuations in state worry, state arousal, and state affect

mood over time. These short manipulation checks were introduced by ‘At this moment...’ and consisted of three domains with each two items: worry (‘... how worried are you?’, ‘... how much do you ruminate?’), arousal (‘... how aroused are you?’, ‘... how tensed are you?’), and affect (‘... how many positive feelings do you feel [e.g., joyful, enthusiastic, active]?’, ‘how many negative feelings do you feel [e.g., angry, sad, anxious]?’). Participants rated each item on a forced-choice visual analogue scale ranging from 0 to 100. Scores of the three domains were aggregated by averaging the respective two items (the item for negative affect was first inverted before averaging). The local ethics committee approved that the study procedure is in accordance with the Declaration of Helsinki (World Medical Association, 2013).

2.3 | Questionnaires

During laboratory assessment, following questionnaires were administered: Trait worry was measured by the Penn State Worry Questionnaire (PSWQ; 16 items, 5-point Likert scale 1–5; $\alpha = .93$; Glöckner-Rist & Rist, 2014; Meyer et al., 1990), anxious arousal by the respective subscale of the Mood and Anxiety Symptom Questionnaire (MASQ-AA; 17 items, 5-point Likert scale 1–5; $\alpha = .87$; Watson & Clark, 1991; Watson et al., 1995), trait anxiety by the respective subscale of the State-Trait-Anxiety Inventory (STAI-T; 20 items, 4-point Likert scale 1–4; $\alpha = .89$; Laux et al., 1981; Spielberger et al., 1970), obsessive-compulsive symptoms by the Obsessive-Compulsive Inventory Revised (OCI-R; 20 items, 5-point Likert scale

0–4; $\alpha = .85$; Foa et al., 2002; Gönner et al., 2007), depression symptoms by the Beck Depression Inventory (BDI-II; 21 items, 4-point Likert scale 0–3; $\alpha = .92$; Beck et al., 1996; Hautzinger et al., 2006), alcohol consumption by the Alcohol Use Disorders Identification Test (AUDIT; 10 items, 5-point Likert scale 0–4; $\alpha = .75$; Babor et al., 2001), and handedness by the modified Edinburgh Handedness Inventory (EHI; 10 items, 5-point Likert scale –10 to +10; $\alpha = .75$; Loffing et al., 2014; Oldfield, 1971).

2.4 | Task

Participants sat in a dimly lit, electrically shielded cabin approx. 24 inches in front of a 19-inch LCD monitor with a resolution 1920 × 1080 pixels and a refresh rate of 120 Hz. A speeded arrowhead version of the flanker task (Eriksen & Eriksen, 1974) with a set of five horizontally aligned arrows (one target, four flankers) was displayed using Presentation Software (Neurobehavioral Systems, Inc., Albany, California). The set of arrows was approx. 6.2° in width and approx. 1.0° in height with an equal number of trials pointing pseudo randomly either into same (<<<<< or >>>>>) or opposite directions (<<><< or >><>>). Each trial included fixation (200–1200 ms), presentation of arrow stimuli (100 ms), and response (max. 800 ms). Participants were instructed to indicate the direction of the center arrow by pressing a key with their respective left or right index finger as quickly and accurately as possible, which was practiced in 20 trials before the first flanker task. Each of the five blocks consisted of 80 trials, which equals 400 trials in total. After each block, participants received a performance feedback asking them to respond faster, irrespective of their actual response times. This procedure was used for three reasons: First, to achieve sufficient error trials in both flanker tasks; second, a reasonable length of the tasks; and third, there is evidence that neural differences between OCD patients and healthy control participants are pronounced under speed conditions (Riesel, Kathmann, et al., 2019).

Accuracy was defined as the percentage of correct responses of the response trials, response times as the time difference between the onset of arrows and the respective correct or incorrect response, and post-error slowing (PES) was quantified using the robust measurement method (i.e., the average response time difference between the last correct trial before an error and the first correct trial after an error; Dutilh et al., 2012).

2.5 | Electrophysiological recording and processing

The setup of recording and processing parameters of the electrophysiological data were mostly as preregistered.

Whenever we deviated from the preregistration, we clearly state which other parameters were chosen. As preregistered, using 61 passive Ag/AgCl-electrodes mounted on a cap with equidistant and concentric electrode sites (EasyCap, Herrsching, Germany) and two 32-channel BrainAmp amplifiers (Brain Products GmbH, Gilching, Germany), EEG signals were recorded with a band-pass filter of 0.01 to 250 Hz and digitized continuously at a sampling rate of 1000 Hz. Recording reference was located between AF3 and Fz, the ground electrode between AF4 and Fz. External electrodes were placed at the left infraorbital site for vertical eye movements and at the neck for the electrocardiogram. Impedances were always kept below 5 k Ω , however, we did not preregister a maximum impedance threshold.

Processing of the EEG data was performed in Brain Vision Analyzer (Brain Products GmbH, Gilching, Germany). First, a band-pass filter with a low cut-off of 0.1 Hz and a high cut-off of 30 Hz (24 dB/oct roll-off) as well as a notch filter of 50 Hz was applied to continuous EEG data. Subsequently, ocular artifacts were corrected by using an independent component analysis (ICA; Jung et al., 2000), whereby relevant components were semi automatically identified and manually checked by visual inspection of the scalp topography, the component activation, and the inspection of the corrected EEG signal. Continuous data were then re-referenced to the average of all scalp electrodes and segmented into response-locked epochs (–500 to 800 ms). Segments containing artifacts (i.e., absolute voltage range exceeding 200 μ V, voltage step exceeding 50 μ V between consecutive data points, or maximum voltage difference of less than 0.5 μ V within 100 ms intervals) were removed. Due to equipment failure ($n = 1$) and low data quality (i.e., more than 25% of all trials containing artifacts; Luck, 2014; $n = 3$) participants were excluded and replaced before data analysis. In the final sample, artifact rejection caused minimal data loss in both the first ($M = 0.99\%$, $SD = 0.02\%$, Max. = 13.00%) and the second flanker ($M = 0.68\%$, $SD = 0.02\%$, Max. = 10.40%). None of participants was falling below a threshold of less than six usable error segments in either flanker task (Olvet & Hajcak, 2009). We did not preregister a fixed baseline interval but specified a visual inspection procedure to identify a baseline that ensures aligned waveforms in the pre-response interval. Consequently, segments were corrected for the baseline interval of –500 to –300 ms and averaged separately for correct and erroneous responses.

The preregistered procedure to determine the electrode at which the ERN and CRN were quantified was also based on visual inspection. At FCz, the signal was maximal after inspecting the grand-averaged waveforms and topographical distributions. As preregistered, quantification was based on peak-to-peak amplitudes, which

is the difference between the most negative peak in the 0–100 ms post-response interval and the most positive peak in a –100 to 0 ms pre-response window. ERPs revealed good psychometric properties for both the ERN ($r_{T0} = .85$; $r_{T1} = .85$) and CRN ($r_{T0} = .96$; $r_{T1} = .92$) as indicated by Spearman-Brown corrected correlations of odd- and even-numbered trials. In order to ensure the robustness of result patterns and to account for the potential influence of methodological choices in ERP research (Klawohn, Meyer, et al., 2020; Sandre et al., 2020), we decided in our preregistration to report results of our main analyses for other commonly used scoring strategies (Tables S1 and S2 in the Supporting Information), including mean amplitude between 0 and 100 ms after response, adaptive mean amplitude around the most negative peak (± 20 ms) in the 0–100 ms post-response interval, and the difference measure (i.e., Δ ERN) using mean amplitude between 0 and 100 ms after correct and erroneous responses.

2.6 | Data analysis

As preregistered, frequentist analyses were conducted using IBM SPSS Statistics, version 25.0 (SPSS, Inc., Chicago) with a significance level of $\alpha = .05$. Differences in demographic (age, education, handedness) and clinical characteristics (all questionnaires) between the three groups were tested by one-way analysis-of-variance (ANOVAs) with group (induction, reduction, control) as between-subject factor. For the manipulation checks, 3×6 mixed-measures ANOVAs with group (induction, reduction, control) as between-subject factor and time (State 1 to 6) as within-subject factor were performed separately for worry, arousal, and affect. Note, that the manipulation check analysis was not specified in the preregistration.

Cross sectional hypotheses on ERPs (ERN and CRN) and behavioral data (accuracy, response times for correct and incorrect response, and PES) were analyzed, as preregistered, using separate multiple linear regression models including the predictors gender, PSWQ, and gender \times PSWQ to investigate the association of worry and ERPs as well as the role of gender as a potential moderator of the relationship between worry and ERPs (Moran et al., 2012; Moser et al., 2016). Exploratively, we also conducted separate independent samples *t*-tests between participants that had any current or lifetime diagnosis and those who were not diagnosed with a disorder to investigate whether these two groups differ in ERN or CRN amplitudes.

Longitudinal hypotheses on ERPs (ERN and CRN) and behavioral data (response times for correct and incorrect response) were analyzed, as preregistered, using separate

$2 \times 2 \times 3$ mixed-measures ANOVAs with time (T0, T1) and response (correct, incorrect) as within-subject factors as well as group (induction, reduction, control) as between-subject factor. Accuracy and PES were tested by 2×3 mixed-measures ANOVAs with time (T0, T1) as within-subject factor and group (induction, reduction, control) as between-subject factor. Greenhouse-Geisser correction was applied if the assumption of sphericity was violated. Follow-up analyses were conducted if results revealed significant interactions. Mirroring the frequentist analyses, we also conducted explorative Bayesian statistical analyses using JASP, version 0.15.0.0. (JASP Team, 2021) allowing the quantification of evidence for the null hypothesis (i.e., the absence of an effect of worry interventions on ERPs; Keyers et al., 2020). Complementary mixed-measures Bayesian ANOVAs were performed with weakly informative priors (r scale fixed effects = 0.5; random effects = 1; covariates = 0.354; van Doorn et al., 2021) resulting in a Bayes factor (BF) that quantifies the evidence for the alternative hypothesis over the null hypothesis (BF_{10}) or for the null hypothesis over the alternative hypothesis ($BF_{01} = 1/BF_{10}$) of a specific model that includes the predictor of interest. For example, the $BF_{10} = 20$ implies that the alternative hypothesis (i.e., there is an effect) is 20 times more likely than the null hypothesis (i.e., the absence of an effect) in light of the data. The Bayes factor included (BF_{Incl}) reflects the inclusion probability of a predictor across all models excluding this specific predictor. Although the Bayes factor is a continuous metric, we refer to a heuristic (Raftery, 1995) when interpreting the BF_{10} and BF_{Incl} : Evidence can be weak ($BF = 1$ –3 or 0.33), positive ($BF = 3$ –20 or 0.33–0.05), strong ($BF = 20$ –150 or 0.05–0.0067), or very strong ($BF > 150$ or 0.0067–0).

3 | RESULTS

3.1 | Demographic and clinical data

Group-specific means and standard deviations of demographic and clinical data can be found in Table 1. As preregistered, we examined potential group differences in these data. However, groups displayed no significant differences regarding demographic and clinical data. In an additional explorative analysis, no differences between groups were found regarding current or lifetime clinical status, $\chi^2(2) = 0.06$, $p = .969$.

3.2 | Manipulation check

Results of the manipulation checks revealed significant differences of participants' mood throughout the study

TABLE 1 Demographical and clinical data of all groups

	Sample (n = 90)		Control (n = 30)		Induction (n = 30)		Reduction (n = 30)		Group Comparison		
	M	SD	M	SD	M	SD	M	SD	F	df	η^2_p
Demographical											
Gender (f/m)	66/24		22/8		22/8		22/8				
Age	23.50	3.12	23.57	3.02	23.20	3.34	23.73	3.07	0.23	2, 87	0.01
Education	12.34	0.60	12.33	0.66	12.47	0.51	12.23	0.63	1.14	2, 87	0.03
Clinical											
PSWQ	45.73	10.93	43.17	9.97	46.83	11.12	47.20	11.54	1.26	2, 87	0.03
MASQ-AA	27.14	8.46	27.77	9.21	25.70	8.04	27.97	8.16	0.66	2, 87	0.01
STAI-T	38.33	8.28	37.33	7.22	37.30	7.73	39.97	9.72	0.89	2, 87	0.02
OCI-R	12.86	8.53	12.77	8.30	14.33	9.36	11.47	7.90	0.85	2, 87	0.02
BDI-II	5.72	6.82	6.40	7.85	4.30	4.56	6.47	7.55	0.98	2, 87	0.02
AUDIT	4.94	3.39	5.00	3.30	4.20	2.89	5.63	3.87	1.36	2, 87	0.03
EHl	81.11	16.21	81.83	15.40	82.00	15.29	79.50	18.21	0.22	2, 87	0.01

Note. Gender (f = female, m = male); age and education in years.
Abbreviations: AUDIT, alcohol use disorders identification test; BDI-II, Beck Depression Inventory II; EHI, Edinburgh Handedness Inventory (Modified); MASQ-AA, Mood and Anxiety Symptom Questionnaire; OCI-R, Obsessive-Compulsive Inventory; PSWQ, Penn State Worry Questionnaire; STAI-T, State-Trait Anxiety Inventory (Trait Subscale).




FIGURE 2 State measurements of state worry, state arousal, and state affect. State measurements tracked fluctuations of state worry (a), state arousal (b), and state affect (c) throughout the study and were assessed before each task. Each scale consisted of two items that were rated on a visual analogue scale. For state worry and state arousal, the potential range was 0 to 100, for state affect it was -50 to 50. Error bars represent one standard error. Asterisks indicate significant differences between groups at each time point: *** $p < .001$. ** $p < .01$. * $p < .05$

(Figure 2). There was no main effect of group, $F(2, 87) = 2.34$, $p = .102$, $\eta_p^2 = 0.05$, but a significant main effect of time, $F(3.60, 312.96) = 16.58$, $p < .001$, $\eta_p^2 = 0.16$, and a significant interaction effect of group \times time, $F(7.19, 312.96) = 10.46$, $p < .001$, $\eta_p^2 = 0.19$, in the ANOVA testing state worry. Follow-up one-way ANOVAs indicated that groups did not differ regarding state worry, neither before the first flanker, $F(2, 87) = 0.04$, $p = .965$, $\eta_p^2 = 0.00$, nor before the intervention, $F(2, 87) = 0.12$, $p = .890$, $\eta_p^2 = 0.00$. However, state worry differed between groups after the intervention, $F(2, 87) = 12.38$, $p < .001$, $\eta_p^2 = 0.22$, indicating that both the induction and the reduction group reported higher levels of state worry compared to the control group (Figure 2). In summary, the induction and reduction group did not differ from the control group regarding state worry before the intervention, but they did so after the intervention during the second flanker and resting state assessment. Consequently, the worry induction successfully increased levels of state worry in the induction group. However, instead of lower levels of state worry in the reduction group, participants reported higher levels of state worry after the emotional expressive writing, which was the opposite pattern of that expected.

Similar patterns were found for state arousal (Figure 2), where there was no main effect of group, $F(2, 86) = 0.73$, $p = .487$, $\eta_p^2 = 0.02$, but a significant effect of time, $F(4.25, 365.22) = 27.28$, $p < .001$, $\eta_p^2 = 0.24$, and group \times time, $F(8.39, 365.22) = 3.59$, $p < .001$, $\eta_p^2 = 0.08$, indicating an increase in arousal as a result of both interventions. State affect also varied as a function of group and time (Figure 2) with no main effect of group, $F(2, 86) = 2.82$, $p = .065$, $\eta_p^2 = 0.06$, but a significant effect of time, $F(3.67, 315.36) = 38.64$, $p < .001$, $\eta_p^2 = 0.31$, and group \times time,

$F(7.33, 315.36) = 5.39$, $p < .001$, $\eta_p^2 = 0.11$. As with state worry and arousal, state positive affect decreased as a result of the worry induction and reduction.¹ Taken together, our interventions not only altered levels of state worry, but also of state arousal and state affect, such that the induction and reduction group did not differ from the control group before the intervention, but did so after the intervention during the second flanker and resting state assessment.

3.3 | Event-related potentials

Group-specific means and standards deviation of the ERN and CRN at T0 and T1 can be found in Table 2.

3.3.1 | Cross-sectional analyses of trait worry and ERPs

The preregistered cross-sectional analyses for the ERN (T0) revealed a significant main effect of gender, $b = 2.65$, $SE = 1.05$, $p = .014$, and an interaction effect of gender \times PSWQ, $b = 0.32$, $SE = 0.10$, $p = .002$, but no main effect of PSWQ, $b = -0.06$, $SE = 0.04$, $p = .221$, $R^2 = .13$, $F(3, 86) = 4.23$, $p = .008$. As indicated by Figure 3, higher levels of trait worry were associated with larger (i.e., more negative) ERN amplitudes in female participants, whereas in male participants, higher levels of trait worry were associated with smaller (i.e., more positive) ERN amplitudes. Regarding the CRN (T0), we observed a main effect of

¹The reason why there is one degree of freedom less in the analyses for arousal and affect was missing data of one participant due to a technical malfunction.

TABLE 2 Group-specific electrophysiological and behavioral data

	Control (<i>n</i> = 30)				Induction (<i>n</i> = 30)				Reduction (<i>n</i> = 30)			
	T0		T1		T0		T1		T0		T1	
	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>	<i>M</i>	<i>SD</i>
ERPs												
ERN (μV)	-7.38	4.33	-8.17	4.67	-7.76	4.72	-8.08	5.25	-6.69	3.30	-6.77	4.07
CRN (μV)	-3.13	2.37	-2.83	2.36	-3.31	2.44	-2.22	2.20	-2.65	1.59	-2.28	1.69
Behavior												
<i>n</i> missing	23.80	54.54	7.03	10.65	15.77	38.63	14.60	42.16	11.70	17.65	6.27	9.33
<i>n</i> correct	329.10	61.54	346.90	44.14	328.63	63.60	327.30	62.18	337.60	34.31	339.73	36.68
<i>n</i> incorrect	47.10	37.62	46.07	40.70	55.60	37.01	58.10	43.32	50.70	29.25	54.00	34.86
Accuracy (%)	87.46	9.51	88.21	10.54	84.72	11.80	84.66	12.51	86.92	7.55	86.27	8.90
RT correct (ms)	437.83	55.41	419.06	51.09	425.65	50.49	397.85	59.59	430.49	53.96	413.82	59.47
RT incorrect (ms)	382.50	58.91	371.18	60.46	365.11	56.58	351.23	66.71	368.09	72.61	362.56	72.41
PES (ms)	39.10	28.45	27.88	30.99	44.56	25.63	25.82	28.42	43.84	24.34	27.47	20.01

Abbreviations: CRN, correct-response negativity; ERN, error-related negativity; ERPs, event-related potentials; PES, post-error slowing; RT, response time.

FIGURE 3 Scatter plots depicting the relationship between trait worry and ERN (a) and CRN (b). CRN, correct-response negativity; ERN, error-related negativity; PSWQ, Penn State Worry Questionnaire. Data points, regression lines, and densities were grouped by gender.





FIGURE 4 Response-locked grand-averaged waveforms of correct and erroneous trials at baseline (T0) for the clinical group with a history of internalizing disorders and the unaffected healthy group (a). Corresponding topographic head maps of the ERN and CRN (b)

gender and PSWQ: Females compared to males showed a higher CRN, $b = 1.20$, $SE = 0.56$, $p = .034$. Furthermore, increasing levels of trait worry were associated with larger CRN amplitudes, $b = -0.05$, $SE = 0.02$, $p = .031$, irrespective of gender since no gender \times PSWQ interaction was observed, $b = 0.09$, $SE = 0.05$, $p = .075$, $R^2 = .11$, $F(3, 86) = 3.44$, $p = .020$.

In addition, we exploratively examined the impact of current or lifetime psychopathology on ERPs: An independent samples t -test compared participants who had any current or lifetime diagnosis of at least one internalizing disorder ($n = 14$) and those who were not diagnosed with a disorder ($n = 69$). Results indicated a larger ERN amplitude in the clinical group ($M = -9.96$, $SD = 4.86$) compared to the healthy group ($M = -6.81$, $SD = 3.68$), $t(81) = 2.76$, $p = .007$, $d = 0.81$. No evidence emerged for CRN associations with lifetime clinical status, $t(81) = 1.45$, $p = .151$, $d = 0.43$ (Figure 4).

3.3.2 | Longitudinal analyses of state worry and ERPs

In the preregistered longitudinal analyses for the ERN and CRN (Figure 5), we found larger amplitudes for erroneous compared to correct responses, $F(1, 87) = 155.49$, $p < .001$, $\eta_p^2 = 0.64$. Further, we found a significant interaction of response \times time, $F(1, 87) = 8.92$, $p = .004$, $\eta_p^2 = 0.09$, driven by a reduction of the CRN at T1, while no evidence emerged for ERN differences between T0 and T1. Importantly, neither a main nor interaction effect including group were observed, suggesting the absence of evidence for worry interventions modulating ERN or CRN. Further, none of the other main or interaction effect reached significance (all $ps > .45$). Detailed statistics are summarized in Table 3.

To further trace the missing intervention effect, we performed non-preregistered Bayesian analyses. These complementary Bayesian analyses yielded very strong




FIGURE 5 Response-locked grand-averaged waveforms of correct and erroneous trials at baseline (T0) and after the intervention (T1) for the worry reduction group (a), passive control group (b), and worry induction group (c). Corresponding topographic head maps of the ERN and CRN for the worry reduction group (d), passive control group (e), and worry induction group (f)

	<i>F</i>	<i>df</i>	η_p^2	<i>p</i>	<i>BF</i> _{Incl.}
Response	155.49	1, 87	0.64	<.001	∞
Response × Time	8.92	1, 87	0.09	.004	0.182
Response × Group	0.46	2, 87	0.01	.632	0.047
Response × Time × Group	0.77	2, 87	0.02	.465	0.000
Time	0.21	1, 87	0.00	.648	0.078
Time × Group	0.80	2, 87	0.02	.453	0.006
Group	0.80	2, 87	0.02	.453	0.084

Note. Bayes factor included (*BF*_{Incl.}), correct-response negativity (CRN), error-related negativity (ERN), group (induction, reduction, control), response (correct, incorrect), time (T0, T1). Significant *p* < .05 printed in bold.

TABLE 3 Results of frequentist and Bayesian ANOVA for ERN and CRN with factors response, time, and group

evidence for response type since all models including response type compared to the null model were more likely given the data ($BF_{10} > 2.61e+39$), with the most likely model encompassing response type alone ($BF_{10} = 8.91e+43$). Furthermore, there was positive to strong evidence against including group, time, or an interaction involving one of them into the model (Table 3) suggesting that neither ERN nor CRN varied as a function of state worry interventions.

Past research suggests that the link between anxiety and larger ERN amplitudes is larger for females (Moser et al., 2016). Therefore, in an explorative approach, we analyzed potential ERN variations of only female participants in our sample ($n = 22$ per group) using a 2×3

mixed-measures ANOVA with time (T0, T1) as a within-subject factor and group (induction, reduction, control) as a between-subject factor. Results indicated no significant effect of group, $F(1, 63) = 0.57$, $p = .567$, $\eta_p^2 = 0.18$, time, $F(1, 63) = 2.92$, $p = .093$, $\eta_p^2 = 0.04$, or time × group, $F(2, 63) = 0.81$, $p = .451$, $\eta_p^2 = 0.03$. Taken together, in the female sample, we could not find evidence for ERN variations from T0 to T1 by our state worry interventions.

3.4 | Behavioral data

Group-specific means and standard deviations of behavioral data at T0 and T1 can be found in Table 2.

3.4.1 | Cross-sectional analyses of trait worry and behavioral data

As preregistered, we also investigated whether behavioral data of the flanker (T0) varied as a function of trait worry and gender. None of the predictor significantly predicted accuracy; neither gender, $b = -0.01$, $SE = 0.03$, $p = .735$, PSWQ, $b = 0.00$, $SE = 0.00$, $p = .855$, nor the interaction of gender \times PSWQ, $b = -0.00$, $SE = 0.00$, $p = .074$, $R^2 = .05$, $F(3, 86) = 1.37$, $p = .259$.

Response times were also not significantly predicted by gender, $b = -24.03$, $SE = 13.75$, $p = .084$ (correct trials), $b = -18.86$, $SE = 16.28$, $p = .250$ (incorrect trials), PSWQ, $b = 1.04$, $SE = 0.59$, $p = .079$ (correct trials), $b = 1.12$, $SE = 0.69$, $p = .111$ (incorrect trials), nor the interaction of gender \times PSWQ, $b = -0.50$, $SE = 1.29$, $p = .646$ (correct trials), $b = 0.78$, $SE = 1.53$, $p = .609$ (incorrect trials), $R^2 = .09$, $F(3, 86) = 2.90$, $p = .040$ (correct trials), $R^2 = .10$, $F(3, 86) = 3.06$, $p = .032$ (incorrect trials).

However, we found an effect of gender on PES: Males compared to females showed less slowing after committing an error, $b = -20.55$, $SE = 6.72$, $p = .003$. No effect on PES emerged for PSWQ, $b = -0.06$, $SE = 0.29$, $p = .847$, or the interaction of gender \times PSWQ, $b = -0.39$, $SE = 0.63$, $p = .537$, $R^2 = .10$, $F(3, 86) = 3.26$, $p = .025$.

When comparing participants with and without a current or lifetime diagnosis of a mental disorder in explorative analyses, we found no differences regarding accuracy, $t(81) = 0.10$, $p = .923$, $d = 0.03$, nor response times, $t(81) = -0.06$, $p = .953$, $d = -0.02$ (correct trials), $t(81) = 0.51$, $p = .611$, $d = 0.15$ (incorrect trials). However, PES was significantly longer in the clinical group compared to the healthy group, $t(81) = -2.19$, $p = .031$, $d = -0.64$.

3.4.2 | Longitudinal analyses of state worry and behavioral data

Following our preregistration, we examined the behavioral data across both sessions and groups. Across groups (induction, reduction, control) and sessions (T0, T1), accuracy was comparable, as indicated by the absence of an effect of group, $F(2, 87) = 0.30$, $p = .446$, $\eta_p^2 = 0.02$, or time, $F(1, 87) = 0.00$, $p = .986$, $\eta_p^2 = 0.00$, in the mixed measures ANOVA. The interaction of group \times time was also not statistically significant, $F(2, 87) = 0.28$, $p = .756$, $\eta_p^2 = 0.01$. For response times, we observed a main effect of response type and session: Participants showed faster responses for incorrect compared with correct trials, $F(1, 87) = 296.79$, $p < .001$, $\eta_p^2 = 0.77$, and were faster at T1 compared to T0, $F(1, 87) = 34.83$, $p < .001$, $\eta_p^2 = 0.29$. No main effect of group and no interaction with group were observed, indicating a comparable training effect in all

groups. In addition, the response \times time interaction was significant, $F(1, 87) = 9.18$, $p = .003$, $\eta_p^2 = 0.10$, reflecting a larger decrease in response times for correct trials. A training effect of time was also found for the PES, where the PES was smaller at the T1, $F(1, 87) = 26.55$, $p < .001$, $\eta_p^2 = 0.24$, independent of group, $F(2, 87) = 0.07$, $p = .930$, $\eta_p^2 = 0.00$, or the interaction of group \times time, $F(2, 87) = 0.52$, $p = .599$, $\eta_p^2 = 0.01$.

4 | DISCUSSION

Previous studies suggested that the link between anxiety and neural correlates of performance monitoring may be driven by worry. However, most of these studies used cross-sectional research designs that preclude causal inferences. The present study aimed at disentangling the effects of trait and state worry on ERN and CRN in a mainly subclinical sample. To this end, we performed cross-sectional as well as longitudinal analyses in a randomized controlled trial. First, $n = 90$ university students completed a flanker task, after which they were randomly assigned to either a worry induction, a worry reduction, or a passive control group. Afterwards, they performed a second flanker task to assess potential alterations of performance related ERPs attributable to the worry interventions. Manipulation checks showed that compared to the control group, state worry increased in the induction group, but also in the reduction group. However, this marked increase in state worry in both groups had no effects on ERN or CRN amplitudes. Across all groups, CRN amplitudes decreased over time, possibly reflecting either training effects or task disengagement. Cross-sectional analyses of the baseline ERPs found larger ERN and CRN amplitudes in females. In addition, higher levels of trait worry were associated with larger CRN amplitudes (irrespective of gender) and larger ERN amplitudes in females only, whereas in males, higher levels of trait worry were associated with smaller ERN amplitudes. Participants with a current or lifetime diagnosis of internalizing disorders showed larger ERN amplitudes compared to participants without a lifetime diagnosis.

In terms of our preregistered hypotheses, we only found partial support for our first hypothesis proposing an association of trait worry with larger ERN amplitudes: Only females showed an increase of ERN with growing levels of trait worry. This finding is in line with meta-analytical evidence suggesting a gender-specific link which was attributed to larger inferences of subvocal rehearsal and gonadal hormones in females on task performance and cognitive control (Moser et al., 2016). Males, in comparison, showed the opposite direction of effects with smaller ERN amplitudes in participants with

increasing trait worry—an unexpected finding that has not been reported before and that warrants further investigation. Nonetheless, we want to emphasize that our sample consisted of only $n = 24$ male participants, limiting the generalizability of male-specific associations. Irrespective of gender, CRN amplitudes were larger in participants with larger trait worry, which might imply that the link to neural correlates of performance monitoring is not specific to error monitoring but translates to performance monitoring in general. This is in line with recent meta-analytical evidence suggesting a small relationship between anxiety and CRN amplitudes (Saunders & Inzlicht, 2020). Our results support this pattern and refine this relationship by pointing to a specific association of trait worry and the CRN. Contradicting previous findings (Fischer et al., 2016; Härpfer, Carsten, Spychalski, et al., 2020; Larson et al., 2011), females showed larger ERN and CRN amplitudes compared to males. As mentioned before, there were only few male participants in our sample limiting the generalizability of the present gender-related results. Future studies targeting research questions of gender-specific effects of performance associated ERPs should recruit more equally distributed samples with large and comparably sized gender groups including participants across the whole worry spectrum, to deliver more informative and reliable evidence for the suggested link. Altogether, our cross-sectional findings support the hypothesized relationship between neural correlates of performance monitoring and worry in females, and suggest that this relationship may not be error-specific.

Because we aimed at disentangling the possible effects of trait and state worry on performance monitoring, we experimentally manipulated state worry by targeted interventions. Because both interventions resulted in a significant increase in state worry, we can only make conclusions about the induction of state worry. No empirical support could be found for our second preregistered hypothesis that higher levels of state worry lead to an increased ERN. This null finding was backed by explorative Bayesian analyses that allow the quantification of evidence for the null hypothesis, yielding positive to strong evidence against an association of state worry and the ERN. In addition, we could also not find evidence for ERN variations due to the state worry interventions in an all-female subsample. Our second hypothesis was derived from the CEMH assuming that the link between error monitoring and worry is the product of compensatory effort of the brain to overcome processing inefficiency due to the workload that worries put on working memory (Moser et al., 2013). However, our results do not support that ERN amplitudes increase when state worry does, as the CEMH predicts.

It is important to note that there is an ongoing debate in the literature whether the ERN is a trait-like neural

risk maker or whether it reflects symptom states. It is also discussed whether the ERN can be altered by interventions (Moser et al., 2013; Proudfit et al., 2013). On the one hand, an error-specific training (Meyer et al., 2020) as well as an attentional bias modification (Klawohn, Hajcak, et al., 2020; Nelson et al., 2015; but see Carlson et al., 2021) successfully reduced ERN amplitudes, suggesting that the ERN can be modulated and might thus be state related to a certain degree. On the other hand, cognitive behavioral therapy decreasing psychopathological symptoms has no effect on the ERN (Gorka et al., 2018; Hajcak et al., 2008; Kujawa et al., 2016; Ladouceur et al., 2018; Riesel et al., 2015), unaffected first-degree relatives and their affected family members show similar aberrant error monitoring (Carrasco et al., 2013; Riesel et al., 2011; Riesel, Klawohn, et al., 2019), and an enhanced ERN is predictive for the onset of anxiety disorders (Meyer et al., 2015). These studies further emphasize the stable, trait-like properties of the ERN but still show that state manipulations that alter error sensitivity also lead to ERN variations.

In light of these previous findings, together with the results of the current study showing ERN variation due to interindividual differences (i.e., trait worry, lifetime internalizing psychopathology) but not intraindividual differences in emotional state (i.e., state worry), we conclude that increased ERN amplitudes are not a consequence or a correlate of worry, but rather may reflect a more stable general disposition (i.e., trait-like) underlying increased anxiety and worry. This conclusion is in line with the assumption that the ERN likely represents a trait-like risk marker or endophenotype for anxiety (Olvet & Hajcak, 2008; Proudfit et al., 2013; Riesel, Klawohn, et al., 2019). However, the fact that the ERN does not seem to be related to state worry does not imply that the ERN per se is insensitive to other emotional states but that the trait-like properties might limit the range of state-associated variability of the ERN. Previous research has found positive affect (Bakic et al., 2014; Larson et al., 2006; Nigbur & Ullsperger, 2020; but see Larson et al., 2013) and negative affect (Pfabigan et al., 2013; Unger et al., 2012; Wiswede, Münte, Goschke, et al., 2009; Wiswede, Münte, & Rüsseler, 2009; but see Larson et al., 2006, 2013) to be associated with elevated ERN amplitudes when manipulating experimental conditions using affective pictures, negative feedback, or mathematical reasoning tasks. Our study design did not aim to directly manipulate positive or negative affect, but as the manipulations checks show, affect was also influenced by the interventions (but to a lesser extent than state worry). The fact that no alterations of the ERN and CRN were observed contradicts earlier findings, but could also be due to an insufficient potency of the intervention. At the same time, the non-specificity

of the effects also illustrates the large overlap between different anxiety-associated emotions.

The interpretation of the present results must be seen in light of certain limitations. Not only was there an unbalanced distribution of gender in our sample, we also recruited mainly participants with subclinical levels of worry. The link between ERN and anxiety was less reliably demonstrated in subclinical samples (Saunders & Inzlicht, 2020) and males (Moser et al., 2016) which might have been the reason why the power of the present study was not large enough to detect small sized effects. However, Bayesian analyses provided positive to strong evidence against an effect of state worry on the ERN.

In addition, we cannot make any statement on the effects of a worry reduction as our intervention to reduce worry levels did not work as intended but instead, increased state worry almost to the same magnitude as the worry induction did. Although there is meta-analytical evidence for emotional expressive writing to improve a broad spectrum of psychological and physiological outcomes (Frattaroli, 2006; Zachariae & O'Toole, 2015), its specific effect on reducing worries has gained less attention. However, there are primary studies showing that expressive writing significantly reduces worries and test anxiety (Goldman et al., 2007; Wolitzky-Taylor & Telch, 2010). Meta-analyses also suggest that effect sizes are larger with increasing duration and number of writing sessions (Reinhold et al., 2018; Travagin et al., 2015). In our study, participants received a very short eight-minute intervention that might have not allowed them to adapt to the exposed worry topics. Another reason might be that we did not pre-select a sample of chronic worriers like Schroder et al. (2018) did. Individuals without chronic worries might not need to reduce or offload their minds from the distracting effects of worries. Therefore, they cannot benefit from such an intervention. However, we have chosen this approach because of the advantage to directly compare effects of a worry induction and reduction within one study.

Regarding the worry induction paradigm, we successfully induced worry, but did not see alterations in the ERN or CRN. This might be due to an absence of state-related influences on these ERPs, but might also be due to an insufficiently potent intervention. In fact, it is still unclear, how potent a worry intervention would have to be in order to alter performance monitoring associated ERPs. A statistically significant increase of state worry must not be equaled with a (clinically) relevant increase. Further, our findings and conclusion are restricted to mainly subclinical populations. Individuals with clinically relevant psychopathologies would be interesting participants for future studies investigating the effects of worry interventions. The effects of the interventions were also not worry-specific but also altered state affect and state arousal. Another difference

to previous studies (Ramirez & Beilock, 2011; Schroder et al., 2018) is that our interventions were not linked to the upcoming testing situation. As a result, the interventions targeted worries that were unrelated to the performance of the flanker task and unrelated worries might have played a subordinate role for participants' performance monitoring at that task. A previous study showed that error monitoring increases with the greater relevance of error commission, such that the ERN is larger when a punishment is related to errors but not when punishment is unrelated (Meyer & Gawlowska, 2017). This might apply to related and unrelated worries as well. Yet, a link to ideographic and task-unrelated worries with personal relevance would imply increased ecological validity of findings and facilitate clinical translation.

In summary, the present study investigated the effects of trait and state worry on performance monitoring associated ERPs (i.e., ERN and CRN) in a mainly subclinical sample. We could only partially replicate an association of trait worry with larger ERN amplitudes: Only females showed larger ERN amplitudes with increasing trait worry. Furthermore, in line with previous studies, we found larger ERN amplitudes in participants with a current or lifetime diagnosis for an internalizing disorder. Concerning the CRN, amplitudes were larger with increasing trait worry (irrespective of gender) but did not differ between participants with or without a lifetime diagnosis of internalizing disorders. Our worry interventions successfully manipulated levels of state worry, but only the worry induction showed effects into the intended direction, whereas the worry reduction also increased state worry. Nevertheless, neither ERN nor CRN amplitudes were altered due to any of the interventions that led to marked increases in state worry. In the face of the current findings, we find tentative support for an association between trait worry and ERN and CRN, with gender being an important moderator. At the same time, we find no evidence for a causal role of state worry influencing the ERN or CRN in our subclinical sample. Instead, consistent with previous findings and assumptions, we demonstrate the stability and independence of the ERN from state worry, as assumed for a trait-like neural risk marker or endophenotype, as which the ERN has been repeatedly conceptualized.

ACKNOWLEDGMENTS

We thank Franziska Ellbracht, Dominika Jeyrani, Moana Langenscheidt, Benedict Lucas, Leonie Osthof, Rica Perkmeyer, Bettina Spitzweck, and Rosann Stocker for their assistance in data acquisition. We also thank the editors and reviewers for their constructive comments and perspectives on our research. Open Access funding enabled and organized by Projekt DEAL.

AUTHOR CONTRIBUTIONS

Kai Härpfer: Conceptualization; data curation; formal analysis; investigation; methodology; project administration; resources; software; visualization; writing – original draft; writing – review and editing. **Hannes Per Carsten:** Conceptualization; investigation; methodology; project administration; validation; writing – review and editing. **Kim Löwisch:** Formal analysis; investigation; validation; visualization; writing – review and editing. **Nele Westermann:** Formal analysis; investigation; validation; visualization; writing – review and editing. **Anja Riesel:** Conceptualization; funding acquisition; methodology; project administration; supervision; writing – review and editing.

CONFLICT OF INTEREST


The authors declare no conflict of interest.

ORCID

Kai Härpfer  <https://orcid.org/0000-0002-9762-2839>

Hannes Per Carsten  <https://orcid.org/0000-0001-7327-4385>

Kim Löwisch  <https://orcid.org/0000-0003-4158-8149>

Nele Westermann  <https://orcid.org/0000-0002-4791-2336>

Anja Riesel  <https://orcid.org/0000-0002-1902-6644>

REFERENCES

- Anokhin, A. P., Golosheykin, S., & Heath, A. C. (2008). Heritability of frontal brain function related to action monitoring. *Psychophysiology*, 45(4), 524–534. <https://doi.org/10.1111/j.1469-8986.2008.00664.x>
- Arch, J. J., & Craske, M. G. (2006). Mechanisms of mindfulness: Emotion regulation following a focused breathing induction. *Behaviour Research and Therapy*, 44(12), 1849–1858. <https://doi.org/10.1016/j.brat.2005.12.007>
- Babor, T. F., Higgins-Biddle, J. C., Saunders, J. B., & Monteiro, M. G. (2001). *AUDIT: The Alcohol Use Disorders Identification Test: Guidelines for use in primary health care* ((2nd ed.) ed.). World Health Organization <https://apps.who.int/iris/handle/10665/67205>
- Baddeley, J. L., & Pennebaker, J. W. (2011). A postdeployment expressive writing intervention for military couples: A randomized controlled trial. *Journal of Traumatic Stress*, 24(5), 581–585. <https://doi.org/10.1002/jts.20679>
- Bakic, J., Jepma, M., De Raedt, R., & Pourtois, G. (2014). Effects of positive mood on probabilistic learning: Behavioral and electrophysiological correlates. *Biological Psychology*, 103, 223–232. <https://doi.org/10.1016/j.biopsycho.2014.09.012>
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Manual for the Beck Depression Inventory-II*. Psychological Corporation.
- Beesdo-Baum, K., Zaudig, M., & Wittchen, H. (2019). *Strukturiertes Klinisches Interview für DSM-5-Störungen—Klinische Version [The Structured Clinical Interview for DSM-5 Disorders—Clinical Version]*. Hogrefe.
- Boehnke, K., Schwartz, S., Stromberg, C., & Sagiv, L. (1998). The structure and dynamics of worry: Theory, measurement, and cross-national replications. *Journal of Personality*, 66(5), 745–782. <https://doi.org/10.1111/1467-6494.00031>
- Borkovec, T. D., & Inz, J. (1990). The nature of worry in generalized anxiety disorder: A predominance of thought activity. *Behaviour Research and Therapy*, 28(2), 153–158. [https://doi.org/10.1016/0005-7967\(90\)90027-G](https://doi.org/10.1016/0005-7967(90)90027-G)
- Botvinick, M. M., Braver, T. S., Barch, D. M., Carter, C. S., & Cohen, J. D. (2001). Conflict monitoring and cognitive control. *Psychological Review*, 108(3), 624–652. <https://doi.org/10.1037/0033-295X.108.3.624>
- Carlson, J. M., Fang, L., & Andrzejewski, J. A. (2021). No change in electrocortical measures of performance monitoring in high trait anxious individuals following multi-session attention bias modification training. *Neuroimage: Reports*, 1(4), 100067. <https://doi.org/10.1016/j.nyirp.2021.100067>
- Carrasco, M., Harbin, S. M., Nienhuis, J. K., Fitzgerald, K. D., Gehring, W. J., & Hanna, G. L. (2013). Increased error-related brain activity in youth with obsessive-compulsive disorder and unaffected siblings. *Depression and Anxiety*, 30(1), 39–46. <https://doi.org/10.1002/da.22035>
- Cavanagh, J. F., & Shackman, A. J. (2015). Frontal midline theta reflects anxiety and cognitive control: Meta-analytic evidence. *Journal of Physiology-Paris*, 109(1), 3–15. <https://doi.org/10.1016/j.jphysparis.2014.04.003>
- Cohen, J. (1992). A power primer. *Psychological Bulletin*, 112(1), 155–159. <https://doi.org/10.1037/0033-2909.112.1.155>
- Dutilh, G., van Ravenzwaaij, D., Nieuwenhuis, S., van der Maas, H. L. J., Forstmann, B. U., & Wagenmakers, E.-J. (2012). How to measure post-error slowing: A confound and a simple solution. *Journal of Mathematical Psychology*, 56(3), 208–216. <https://doi.org/10.1016/j.jmp.2012.04.001>
- Endrass, T., Riesel, A., Kathmann, N., & Buhlmann, U. (2014). Performance monitoring in obsessive-compulsive disorder and social anxiety disorder. *Journal of Abnormal Psychology*, 123(4), 705–714. <https://doi.org/10.1037/abn0000012>
- Eriksen, B. A., & Eriksen, C. W. (1974). Effects of noise letters upon the identification of a target letter in a nonsearch task. *Perception & Psychophysics*, 16(1), 143–149. <https://doi.org/10.3758/BF03203267>
- Falkenstein, M., Hohnsbein, J., Hoormann, J., & Blanke, L. (1991). Effects of crossmodal divided attention on late ERP components: II Error processing in choice reaction tasks. *Electroencephalography & Clinical Neurophysiology*, 78(6), 447–455. [https://doi.org/10.1016/0013-4694\(91\)90062-9](https://doi.org/10.1016/0013-4694(91)90062-9)
- Faul, F., Erdfelder, E., Buchner, A., & Lang, A.-G. (2009). Statistical power analyses using G*Power 3.1: Tests for correlation and regression analyses. *Behavior Research Methods*, 41, 1149–1160. <http://gpower.hhu.de>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*Power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods*, 39, 175–191. <http://gpower.hhu.de>
- First, M., Williams, J., Karg, R., & Spitzer, R. (2016). *Structured clinical interview for DSM-5 disorders, clinician version (SCID-5-CV)*. American Psychiatric Association.
- Fischer, A. G., Danielmeier, C., Villringer, A., Klein, T. A., & Ullsperger, M. (2016). Gender influences on brain responses to errors and post-error adjustments. *Scientific Reports*, 6(1), 24435. <https://doi.org/10.1038/srep24435>
- Fisher, A. J., & Newman, M. G. (2013). Heart rate and autonomic response to stress after experimental induction of worry versus

- relaxation in healthy, high-worry, and generalized anxiety disorder individuals. *Biological Psychology*, 93(1), 65–74. <https://doi.org/10.1016/j.biopsycho.2013.01.012>
- Foa, E. B., Huppert, J. D., Leiberg, S., Langner, R., Kichic, R., Hajcak, G., & Salkovskis, P. M. (2002). The Obsessive-Compulsive Inventory: Development and validation of a short version. *Psychological Assessment*, 14(4), 485–496. <https://doi.org/10.1037/1040-3590.14.4.485>
- Frattaroli, J. (2006). Experimental disclosure and its moderators: A meta-analysis. *Psychological Bulletin*, 132(6), 823–865. <https://doi.org/10.1037/0033-2909.132.6.823>
- Gehring, W. J., Goss, B., Coles, M. G., Meyer, D. E., & Donchin, E. (1993). A neural system for error detection and compensation. *Psychological Science*, 4(6), 385–390. <https://doi.org/10.1111/j.1467-9280.1993.tb00586.x>
- Glöckner-Rist, A., & Rist, F. (2014). Deutsche Version des Penn State Worry Questionnaire (PSWQ-d) [German version of the Penn State Worry Questionnaire]. *Zusammenstellung sozialwissenschaftlicher Items und Skalen*, 1–26. <https://doi.org/10.6102/zis219>
- Goldman, N., Dugas, M. J., Sexton, K. A., & Gervais, N. J. (2007). The impact of written exposure on worry: A preliminary investigation. *Behavior Modification*, 31(4), 512–538. <https://doi.org/10.1177/0145445506298651>
- Gönner, S., Leonhart, R., & Ecker, W. (2007). Das Zwangsinventar OCI-R - Die deutsche Version des Obsessive-Compulsive Inventory-Revised [The German Version of the Obsessive-Compulsive Inventory-Revised]. *Psychotherapie Psychosomatik Medizinische Psychologie*, 57(09/10), 395–404. <https://doi.org/10.1055/s-2007-970894>
- Gorka, S. M., Burkhouse, K. L., Klumpp, H., Kennedy, A. E., Afshar, K., Francis, J., Ajilore, O., Mariouw, S., Craske, M. G., Langenecker, S., Shankman, S. A., & Phan, K. L. (2018). Error-related brain activity as a treatment moderator and index of symptom change during cognitive-behavioral therapy or selective serotonin reuptake inhibitors. *Neuropsychopharmacology*, 43(6), 1355–1363. <https://doi.org/10.1038/npp.2017.289>
- Gortner, E.-M., Rude, S. S., & Pennebaker, J. W. (2006). Benefits of expressive writing in lowering rumination and depressive symptoms. *Behavior Therapy*, 37(3), 292–303. <https://doi.org/10.1016/j.beth.2006.01.004>
- Gottesman, I. I., & Gould, T. D. (2003). The endophenotype concept in psychiatry: Etymology and strategic intentions. *The American Journal of Psychiatry*, 160(4), 636–645. <https://doi.org/10.1176/appi.ajp.160.4.636>
- Hajcak, G., Franklin, M. E., Foa, E. B., & Simons, R. F. (2008). Increased error-related brain activity in pediatric obsessive-compulsive disorder before and after treatment. *American Journal of Psychiatry*, 165(1), 116–123. <https://doi.org/10.1176/appi.ajp.2007.07010143>
- Hajcak, G., Klawohn, J., & Meyer, A. (2019). The utility of event-related potentials in clinical psychology. *Annual Review of Clinical Psychology*, 15(1), 71–95. <https://doi.org/10.1146/annurev-clinpsy-050718-095457>
- Hajcak, G., McDonald, N., & Simons, R. F. (2003). Anxiety and error-related brain activity. *Biological Psychology*, 64(1-2), 77–90. [https://doi.org/10.1016/S0301-0511\(03\)00103-0](https://doi.org/10.1016/S0301-0511(03)00103-0)
- Härpfer, K., Carsten, H. P., & Riesel, A. (2020, August 31). *The modulating effect of trait and state worry on the error-related negativity (ERN)*. <https://osf.io/j8k6z>
- Härpfer, K., Carsten, H. P., Spychalski, D., Kathmann, N., & Riesel, A. (2020). Were we erring? The impact of worry and arousal on error-related negativity in a non-clinical sample. *Psychophysiology*, 57(11), e13661. <https://doi.org/10.1111/psyp.13661>
- Hautzinger, M., Keller, F., & Kühner, C. (2006). *Beck Depressions-Inventar (BDI-II)*. Harcourt Test Services.
- Holroyd, C. B., & Coles, M. G. H. (2002). The neural basis of human error processing: Reinforcement learning, dopamine, and the error-related negativity. *Psychological Review*, 109(4), 679–709. <https://doi.org/10.1037/0033-295X.109.4.679>
- JASP Team. (2021). *JASP (Version 0.15.0.0) [Computer software]*. <https://jasp-stats.org/>
- Jung, T.-P., Makeig, S., Humphries, C., Lee, T.-W., McKeown, M. J., Iragui, V., & Sejnowski, T. J. (2000). Removing electroencephalographic artifacts by blind source separation. *Psychophysiology*, 37(2), 163–178. <https://doi.org/10.1111/1469-8986.3720163>
- Keyesers, C., Gazzola, V., & Wagenmakers, E.-J. (2020). Using Bayes factor hypothesis testing in neuroscience to establish evidence of absence. *Nature Neuroscience*, 23(7), 788–799. <https://doi.org/10.1038/s41593-020-0660-4>
- Klawohn, J., Hajcak, G., Amir, N., Kathmann, N., & Riesel, A. (2020). Application of attentional bias modification training to modulate hyperactive error-monitoring in OCD. *International Journal of Psychophysiology*, 156, 79–86. <https://doi.org/10.1016/j.ijpsycho.2020.07.005>
- Klawohn, J., Meyer, A., Weinberg, A., & Hajcak, G. (2020). Methodological choices in event-related potential (ERP) research and their impact on internal consistency reliability and individual differences: An examination of the error-related negativity (ERN) and anxiety. *Journal of Abnormal Psychology*, 129(1), 29–37. <https://doi.org/10.1037/abn0000458>
- Kujawa, A., Weinberg, A., Bunford, N., Fitzgerald, K. D., Hanna, G. L., Monk, C. S., Kennedy, A. E., Klumpp, H., Hajcak, G., & Phan, K. L. (2016). Error-related brain activity in youth and young adults before and after treatment for generalized or social anxiety disorder. *Progress in Neuro-Psychopharmacology and Biological Psychiatry*, 71, 162–168. <https://doi.org/10.1016/j.pnpbp.2016.07.010>
- Ladouceur, C. D., Tan, P. Z., Sharma, V., Bylsma, L. M., Silk, J. S., Siegle, G. J., Forbes, E. E., McMakin, D. L., Dahl, R. E., Kendall, P. C., Mannarino, A., & Ryan, N. D. (2018). Error-related brain activity in pediatric anxiety disorders remains elevated following individual therapy: A randomized clinical trial. *Journal of Child Psychology and Psychiatry*, 59(11), 1152–1161. <https://doi.org/10.1111/jcpp.12900>
- Larson, M. J., Gray, A. C., Clayson, P. E., Jones, R., & Kirwan, C. B. (2013). What are the influences of orthogonally-manipulated valence and arousal on performance monitoring processes? The effects of affective state. *International Journal of Psychophysiology*, 87(3), 327–339. <https://doi.org/10.1016/j.ijpsycho.2013.01.005>
- Larson, M. J., Perlstein, W. M., Stigge-Kaufman, D., Kelly, K. G., & Dotson, V. M. (2006). Affective context-induced modulation of the error-related negativity. *NeuroReport*, 17(3). <https://doi.org/10.1097/01.wnr.0000199461.01542.db>
- Larson, M. J., South, M., & Clayson, P. E. (2011). Sex differences in error-related performance monitoring. *NeuroReport*, 22(1), 44–48. <https://doi.org/10.1097/WNR.0b013e3283427403>

- Laux, L., Glanzmann, P., Schaffner, P., & Spielberger, C. D. (1981). *Das State-Trait-Angstinventar (STAI) [The State-Trait Anxiety Inventory (STAI)]*. Beltz.
- Lin, Y., Moran, T. P., Schroder, H. S., & Moser, J. S. (2015). The role of hand of error and stimulus orientation in the relationship between worry and error-related brain activity: Implications for theory and practice. *Psychophysiology*, 52(10), 1281–1292. <https://doi.org/10.1111/psyp.12470>
- Loffing, F., Sölter, F., & Hagemann, N. (2014). Left preference for sport tasks does not necessarily indicate left-handedness: Sport-specific lateral preferences, relationship with handedness and implications for laterality research in behavioural sciences. *PLOS ONE*, 9(8), e105800. <https://doi.org/10.1371/journal.pone.0105800>
- Luck, S. J. (2014). *An introduction to the event-related potential technique* ((2nd ed.). ed.). MIT Press.
- Luijten, M., Machielsen, M. W. J., Veltman, D. J., Hester, R., de Haan, L., & Franken, I. H. A. (2014). Systematic review of ERP and fMRI studies investigating inhibitory control and error processing in people with substance dependence and behavioural addictions. *Journal of Psychiatry & Neuroscience*, 39(3), 149–169. <https://doi.org/10.1503/jpn.130052>
- Lutz, M. C., Kok, R., Verveer, I., Malbec, M., Koot, S., van Lier, P. A. C., & Franken, I. H. A. (2021). Diminished error-related negativity and error positivity in children and adults with externalizing problems and disorders: A meta-analysis on error processing. *Journal of Psychiatry & Neuroscience*, 46(6), E615. <https://doi.org/10.1503/jpn.200031>
- Lyonfields, J. D., Borkovec, T. D., & Thayer, J. F. (1995). Vagal tone in generalized anxiety disorder and the effects of aversive imagery and worrisome thinking. *Behavior Therapy*, 26(3), 457–466. [https://doi.org/10.1016/S0005-7894\(05\)80094-2](https://doi.org/10.1016/S0005-7894(05)80094-2)
- Manoach, D. S., & Agam, Y. (2013). Neural markers of errors as endophenotypes in neuropsychiatric disorders. *Frontiers in Human Neuroscience*, 7, 350. <https://doi.org/10.3389/fnhum.2013.00350>
- Mathews, C. A., Perez, V. B., Delucchi, K. L., & Mathalon, D. H. (2012). Error-related negativity in individuals with obsessive-compulsive symptoms: Toward an understanding of hoarding behaviors. *Biological Psychology*, 89(2), 487–494. <https://doi.org/10.1016/j.biopsycho.2011.12.018>
- McLaughlin, K. A., Borkovec, T. D., & Sibrava, N. J. (2007). The effects of worry and rumination on affect states and cognitive activity. *Behavior Therapy*, 38(1), 23–38. <https://doi.org/10.1016/j.beth.2006.03.003>
- Meyer, A., & Gawlowska, M. (2017). Evidence for specificity of the impact of punishment on error-related brain activity in high versus low trait anxious individuals. *International Journal of Psychophysiology*, 120, 157–163. <https://doi.org/10.1016/j.ijpsycho.2017.08.001>
- Meyer, A., Gibby, B., Wisemann, K., Klawohn, J., Hajcak, G., & Schmidt, N. B. (2020). A brief, computerized intervention targeting error sensitivity reduces the error-related negativity. *Cognitive, Affective, & Behavioral Neuroscience*, 20(1), 172–180. <https://doi.org/10.3758/s13415-019-00760-w>
- Meyer, A., Hajcak, G., Torpey-Newman, D. C., Kujawa, A., & Klein, D. N. (2015). Enhanced error-related brain activity in children predicts the onset of anxiety disorders between the ages of 6 and 9. *Journal of Abnormal Psychology*, 124(2), 266–274. <https://doi.org/10.1037/abn0000044>
- Meyer, A., Weinberg, A., Klein, D. N., & Hajcak, G. (2012). The development of the error-related negativity (ERN) and its relationship with anxiety: Evidence from 8 to 13 year-olds. *Developmental Cognitive Neuroscience*, 2(1), 152–161. <https://doi.org/10.1016/j.dcn.2011.09.005>
- Meyer, T. J., Miller, M. L., Metzger, R. L., & Borkovec, T. D. (1990). Development and validation of the Penn State Worry Questionnaire. *Behaviour Research and Therapy*, 28(6), 487–495. [https://doi.org/10.1016/0005-7967\(90\)90135-6](https://doi.org/10.1016/0005-7967(90)90135-6)
- Moran, T. P., Taylor, D., & Moser, J. S. (2012). Sex moderates the relationship between worry and performance monitoring brain activity in undergraduates. *International Journal of Psychophysiology*, 85(2), 188–194. <https://doi.org/10.1016/j.ijpsycho.2012.05.005>
- Moser, J. S., Moran, T. P., & Jendrusina, A. A. (2012). Parsing relationships between dimensions of anxiety and action monitoring brain potentials in female undergraduates. *Psychophysiology*, 49(1), 3–10. <https://doi.org/10.1111/j.1469-8986.2011.01279.x>
- Moser, J. S., Moran, T. P., Kneip, C., Schroder, H. S., & Larson, M. J. (2016). Sex moderates the association between symptoms of anxiety, but not obsessive compulsive disorder, and error-monitoring brain activity: A meta-analytic review. *Psychophysiology*, 53(1), 21–29. <https://doi.org/10.1111/psyp.12509>
- Moser, J. S., Moran, T. P., Schroder, H. S., Donnellan, M. B., & Yeung, N. (2013). On the relationship between anxiety and error monitoring: A meta-analysis and conceptual framework. *Frontiers in Human Neuroscience*, 7, 466. <https://doi.org/10.3389/fnhum.2013.00466>
- Nelson, B. D., Jackson, F., Amir, N., & Hajcak, G. (2015). Single-session attention bias modification and error-related brain activity. *Cognitive, Affective, & Behavioral Neuroscience*, 15(4), 776–786. <https://doi.org/10.3758/s13415-015-0365-4>
- Nigbur, R., & Ullsperger, M. (2020). Funny kittens: Positive mood induced via short video-clips affects error processing but not conflict control. *International Journal of Psychophysiology*, 147, 147–155. <https://doi.org/10.1016/j.ijpsycho.2019.11.007>
- Oathes, D. J., Ray, W. J., Yamasaki, A. S., Borkovec, T. D., Castonguay, L. G., Newman, M. G., & Nitschke, J. (2008). Worry, generalized anxiety disorder, and emotion: Evidence from the EEG gamma band. *Biological Psychology*, 79(2), 165–170. <https://doi.org/10.1016/j.biopsycho.2008.04.005>
- Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, 9(1), 97–113. [https://doi.org/10.1016/0028-3932\(71\)90067-4](https://doi.org/10.1016/0028-3932(71)90067-4)
- Olivet, D. M., & Hajcak, G. (2008). The error-related negativity (ERN) and psychopathology: Toward an endophenotype. *Clinical Psychology Review*, 28(8), 1343–1354. <https://doi.org/10.1016/j.cpr.2008.07.003>
- Olivet, D. M., & Hajcak, G. (2009). The stability of error-related brain activity with increasing trials. *Psychophysiology*, 46(5), 957–961. <https://doi.org/10.1111/j.1469-8986.2009.00848.x>
- Pasion, R., & Barbosa, F. (2019). ERN as a transdiagnostic marker of the internalizing-externalizing spectrum: A dissociable meta-analytic effect. *Neuroscience and Biobehavioral Reviews*, 103, 133–149. <https://doi.org/10.1016/j.neubiorev.2019.06.013>
- Pennebaker, J. W., & Beall, S. K. (1986). Confronting a traumatic event: Toward an understanding of inhibition and disease. *Journal of Abnormal Psychology*, 95(3), 274–281. <https://doi.org/10.1037/0021-843x.95.3.274>

- Pennebaker, J. W., & Francis, M. E. (1996). Cognitive, emotional, and language processes in disclosure. *Cognition and Emotion*, 10(6), 601–626. <https://doi.org/10.1080/026999396380079>
- Pfabigan, D. M., Pintzinger, N. M., Siedek, D. R., Lamm, C., Derntl, B., & Sailer, U. (2013). Feelings of helplessness increase ERN amplitudes in healthy individuals. *Neuropsychologia*, 51(4), 613–621. <https://doi.org/10.1016/j.neuropsychologia.2012.12.008>
- Proudfit, G. H., Inzlicht, M., & Mennin, D. S. (2013). Anxiety and error monitoring: The importance of motivation and emotion. *Frontiers in Human Neuroscience*, 7, 636. <https://doi.org/10.3389/fnhum.2013.00636>
- Raftery, A. E. (1995). Bayesian Model Selection in Social Research. *Sociological Methodology*, 25, 111–163. <https://doi.org/10.2307/271063>
- Ramirez, G., & Beilock, S. L. (2011). Writing about testing worries boosts exam performance in the classroom. *Science*, 331(6014), 211–213. <https://doi.org/10.1126/science.1199427>
- Ray, W. J., Molnar, C., Aikins, D., Yamasaki, A., Newman, M. G., Castonguay, L., & Borkovec, T. D. (2009). Startle response in generalized anxiety disorder. *Depression and Anxiety*, 26(2), 147–154. <https://doi.org/10.1002/da.20479>
- Reinhold, M., Bürkner, P.-C., & Holling, H. (2018). Effects of expressive writing on depressive symptoms—A meta-analysis. *Clinical Psychology: Science and Practice*, 25(1), e12224. <https://doi.org/10.1111/cpsp.12224>
- Riesel, A. (2019). The erring brain: Error-related negativity as an endophenotype for OCD—A review and meta-analysis. *Psychophysiology*, 56(4), e13348. <https://doi.org/10.1111/psyp.13348>
- Riesel, A., Endrass, T., Auerbach, L. A., & Kathmann, N. (2015). Overactive performance monitoring as an endophenotype for obsessive-compulsive disorder: Evidence from a treatment study. *American Journal of Psychiatry*, 172(7), 665–673. <https://doi.org/10.1176/appi.ajp.2014.14070886>
- Riesel, A., Endrass, T., Kaufmann, C., & Kathmann, N. (2011). Overactive error-related brain activity as a candidate endophenotype for obsessive-compulsive disorder: Evidence from unaffected first-degree relatives. *The American Journal of Psychiatry*, 168(3), 317–324. <https://doi.org/10.1176/appi.ajp.2010.10030416>
- Riesel, A., Goldhahn, S., & Kathmann, N. (2017). Hyperactive performance monitoring as a transdiagnostic marker: Results from health anxiety in comparison to obsessive-compulsive disorder. *Neuropsychologia*, 96, 1–8. <https://doi.org/10.1016/j.neuropsychologia.2016.12.029>
- Riesel, A., Kathmann, N., & Klawohn, J. (2019). Flexibility of error-monitoring in obsessive-compulsive disorder under speed and accuracy instructions. *Journal of Abnormal Psychology*, 128(7), 671–677. <https://doi.org/10.1037/abn0000463>
- Riesel, A., Klawohn, J., Grützmann, R., Kaufmann, C., Heinzl, S., Bey, K., Lennertz, L., Wagner, M., & Kathmann, N. (2019). Error-related brain activity as a transdiagnostic endophenotype for obsessive-compulsive disorder, anxiety and substance use disorder. *Psychological Medicine*, 49(7), 1–11. <https://doi.org/10.1017/S0033291719000199>
- Riesel, A., Weinberg, A., Endrass, T., Kathmann, N., & Hajcak, G. (2012). Punishment has a lasting impact on error-related brain activity. *Psychophysiology*, 49(2), 239–247. <https://doi.org/10.1111/j.1469-8986.2011.01298.x>
- Ruscio, A. M., & Borkovec, T. D. (2004). Experience and appraisal of worry among high worriers with and without generalized anxiety disorder. *Behaviour Research and Therapy*, 42(12), 1469–1482. <https://doi.org/10.1016/j.brat.2003.10.007>
- Sandre, A., Banica, I., Riesel, A., Flake, J., Klawohn, J., & Weinberg, A. (2020). Comparing the effects of different methodological decisions on the error-related negativity and its association with behaviour and gender. *International Journal of Psychophysiology*, 156, 18–39. <https://doi.org/10.1016/j.ijpsycho.2020.06.016>
- Saunders, B., & Inzlicht, M. (2020). Assessing and adjusting for publication bias in the relationship between anxiety and the error-related negativity. *International Journal of Psychophysiology*, 155, 87–98. <https://doi.org/10.1016/j.ijpsycho.2020.05.008>
- Sayer, N. A., Noorbaloochi, S., Frazier, P. A., Pennebaker, J. W., Orazem, R. J., Schnurr, P. P., Murdoch, M., Carlson, K. F., Gravely, A., & Litz, B. T. (2015). Randomized controlled trial of online expressive writing to address readjustment difficulties among U.S. Afghanistan and Iraq war veterans. *Journal of Traumatic Stress*, 28(5), 381–390. <https://doi.org/10.1002/jts.22047>
- Schroder, H. S., Moran, T. P., & Moser, J. S. (2018). The effect of expressive writing on the error-related negativity among individuals with chronic worry. *Psychophysiology*, 55(2), e12990. <https://doi.org/10.1111/psyp.12990>
- Shiels, K., & Hawk, L. W. (2010). Self-regulation in ADHD: The role of error processing. *Clinical Psychology Review*, 30(8), 951–961. <https://doi.org/10.1016/j.cpr.2010.06.010>
- Simons, R. F. (2010). The way of our errors: Theme and variations. *Psychophysiology*, 47(1), 1–14. <https://doi.org/10.1111/j.1469-8986.2009.00929.x>
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the State-Trait Anxiety Inventory*. Consulting Psychologists Press.
- Suor, J. H., Calentino, A. E., Granros, M., & Burkhouse, K. L. (2021). Maternal error-related negativity relationship with offspring error-related negativity and negative parenting styles: A novel model of internalizing psychopathology risk. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*. <https://doi.org/10.1016/j.bpsc.2021.10.007>
- Thayer, J. F., Friedman, B. H., & Borkovec, T. D. (1996). Autonomic characteristics of generalized anxiety disorder and worry. *Biological Psychiatry*, 39(4), 255–266. [https://doi.org/10.1016/0006-3223\(95\)00136-0](https://doi.org/10.1016/0006-3223(95)00136-0)
- Travagin, G., Margola, D., & Revenson, T. A. (2015). How effective are expressive writing interventions for adolescents? A meta-analytic review. *Clinical Psychology Review*, 36, 42–55. <https://doi.org/10.1016/j.cpr.2015.01.003>
- Unger, K., Kray, J., & Mecklinger, A. (2012). Worse than feared? Failure induction modulates the electrophysiological signature of error monitoring during subsequent learning. *Cognitive, Affective, & Behavioral Neuroscience*, 12(1), 34–51. <https://doi.org/10.3758/s13415-011-0061-y>
- van Doorn, J., van den Bergh, D., Böhm, U., Dablander, F., Derks, K., Draws, T., Etz, A., Evans, N. J., Gronau, Q. F., Haaf, J. M., Hinne, M., Kucharský, Š., Ly, A., Marsman, M., Matzke, D., Gupta, A. R. K. N., Sarafoglou, A., Stefan, A., Voelkel, J. G., & Wagenmakers, E.-J. (2021). The JASP guidelines for conducting and reporting a Bayesian analysis. *Psychonomic Bulletin & Review*, 28(3), 813–826. <https://doi.org/10.3758/s13423-020-01798-5>

- Vasey, M. W., & Borkovec, T. D. (1992). A catastrophizing assessment of worrisome thoughts. *Cognitive Therapy and Research*, 16(5), 505–520. <https://doi.org/10.1007/BF01175138>
- Verkuil, B., Brosschot, J. F., Putman, P., & Thayer, J. F. (2009). Interacting effects of worry and anxiety on attentional disengagement from threat. *Behaviour Research and Therapy*, 47(2), 146–152. <https://doi.org/10.1016/j.brat.2008.11.003>
- Vidal, F., Burle, B., Bonnet, M., Grapperon, J., & Hasbroucq, T. (2003). Error negativity on correct trials: A reexamination of available data. *Biological Psychology*, 64(3), 265–282. [https://doi.org/10.1016/S0301-0511\(03\)00097-8](https://doi.org/10.1016/S0301-0511(03)00097-8)
- Vidal, F., Hasbroucq, T., Grapperon, J., & Bonnet, M. (2000). Is the 'error negativity' specific to errors? *Biological Psychology*, 51(2–3), 109–128. [https://doi.org/10.1016/S0301-0511\(99\)00032-0](https://doi.org/10.1016/S0301-0511(99)00032-0)
- Watson, D., & Clark, L. A. (1991). *The Mood and Anxiety Symptom Questionnaire*. University of Iowa.
- Watson, D., Weber, K., Assenheimer, J. S., Clark, L. A., Strauss, M. E., & McCormick, R. A. (1995). Testing a tripartite model: I Evaluating the convergent and discriminant validity of anxiety and depression symptom scales. *Journal of Abnormal Psychology*, 104(1), 3–14. <https://doi.org/10.1037/0021-843X.104.1.3>
- Weinberg, A., Olvet, D. M., & Hajcak, G. (2010). Increased error-related brain activity in generalized anxiety disorder. *Biological Psychology*, 85(3), 472–480. <https://doi.org/10.1016/j.biopsycho.2010.09.011>
- Weinberg, A., Riesel, A., & Hajcak, G. (2012). Integrating multiple perspectives on error-related brain activity: The ERN as a neural indicator of trait defensive reactivity. *Motivation and Emotion*, 36(1), 84–100. <https://doi.org/10.1007/s11031-011-9269-y>
- Wiswede, D., Münte, T. F., Goschke, T., & Rüsseler, J. (2009). Modulation of the error-related negativity by induction of short-term negative affect. *Neuropsychologia*, 47(1), 83–90. <https://doi.org/10.1016/j.neuropsychologia.2008.08.016>
- Wiswede, D., Münte, T. F., & Rüsseler, J. (2009). Negative affect induced by derogatory verbal feedback modulates the neural signature of error detection. *Social Cognitive and Affective Neuroscience*, 4(3), 227–237. <https://doi.org/10.1093/scan/nsp015>
- Wolitzky-Taylor, K. B., & Telch, M. J. (2010). Efficacy of self-administered treatments for pathological academic worry: A randomized controlled trial. *Behaviour Research and Therapy*, 48(9), 840–850. <https://doi.org/10.1016/j.brat.2010.03.019>
- World Medical Association. (2013). World Medical Association declaration of Helsinki: Ethical principles for medical research involving human subjects. *JAMA*, 310(20), 2191–2194. <https://doi.org/10.1001/jama.2013.281053>
- Yeung, N., Botvinick, M. M., & Cohen, J. D. (2004). The neural basis of error detection: Conflict monitoring and the error-related negativity. *Psychological Review*, 111(4), 931–959. <https://doi.org/10.1037/0033-295X.111.4.931>
- Zachariae, R., & O'Toole, M. S. (2015). The effect of expressive writing intervention on psychological and physical health outcomes in cancer patients—A systematic review and meta-analysis. *Psycho-Oncology*, 24(11), 1349–1359. <https://doi.org/10.1002/pon.3802>

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

FIGURE S1 Output with specifications of the a priori sample size calculation using G*Power, version 3.1.9.7

TABLE S1 Effects of gender, PSWQ, and Gender × PSWQ on the ERN and CRN using different scoring approaches

TABLE S2 Results of frequentist ANOVA for ERN and CRN varying as function of time and group using different scoring approaches

How to cite this article: Härpfer, K., Carsten, H. P., Löwisch, K., Westermann, N. & Riesel, A. (2022). Disentangling the effects of trait and state worry on error-related brain activity: Results from a randomized controlled trial using worry manipulations. *Psychophysiology*, 59, e14055. <https://doi.org/10.1111/psyp.14055>